

Table SII. Proportion of patients with alopecia above the threshold ($T > 65$) for the different Minnesota Multiphasic Personality Inventory (MMPI) scales, by gender, duration of disease, and self-reported stress

	Gender		Duration of disease			Self-reported stress	
	Male $n=33$ %	Female $n=40$ %	<6 ($n=36$) %	6–11 ($n=15$) %	>12 ($n=22$) %	Yes %	No %
Lie	3.0	7.5	8.3	0	0	2.7	8.3
Frequency	9.1	5.0	5.6	23.1	0**	10.8	2.8
Correction	3.0	2.5	2.8	0	4.5	5.4	0
Hypochondriasis	9.1	12.5	11.1	23.1	4.5	13.5	8.3
Depression	21.2	7.5	13.9	23.1	9.1	16.2	11.1
Hysteria	12.1	10.0	11.1	30.8	0**	13.5	8.3
Psychopathic deviance	21.2	5.0*	5.6	30.8	13.6*	10.8	13.9
Masculinity/Femininity	9.1	0*	5.6	0	4.5	2.7	5.6
Paranoia	6.1	10.0	2.8	30.8	4.5***	8.1	8.3
Psychasthenia	12.1	10.0	8.3	30.8	4.5**	13.5	8.3
Schizophrenia	6.1	10.0	2.8	30.8	4.5***	13.5	2.8*
Hypomania	9.1	10.0	13.9	15.4	0	8.1	11.1
Social introversion	6.1	7.5	11.1	0	4.5	5.4	8.3
Anxiety	18.2	10.0	11.1	30.8	9.1	18.9	8.3
Fears	12.1	2.5	5.6	15.4	4.5	8.1	5.6
Obsessiveness	6.1	0	2.8	0	4.5	2.7	2.8
Depression	15.1	10.0	11.1	30.8	4.5*	18.9	5.6*
Health concerns	15.1	17.5	19.4	15.4	13.6	18.9	13.9
Bizarre thoughts	6.1	10.0	5.6	23.1	4.5	8.1	8.3
Anger	9.1	5.0	5.6	23.1	0**	5.4	8.3
Cynicism	0	2.5	2.8	0	0	0	2.8
Antisocial practices	9.1	0*	2.8	15.4	0*	2.7	5.6
Type A behaviour	3.0	0	2.8	0	0	2.7	0
Low self-esteem	3.0	0	2.8	0	0	2.7	0
Social discomfort	6.1	2.5	5.6	0	4.5	5.4	2.8
Family problems	15.1	2.5*	2.8	23.1	9.1*	10.8	5.6
Work interference	6.1	0	5.6	0	0	5.4	0
Negative treatment	0	2.5	2.8	0	0		

Significance tested using Fisher's exact test is shown in bold; * $p < 0.10$; ** $p < 0.05$; *** $p < 0.01$. Raw scores on the scales are transformed into a standardized metric known as T-score (mean 50, and standard deviation 10).