

Table SII. Prevalence of suicidal ideation in itch, pain, family income and ethnicity among 1,621 boys and 2,061 girls in Oslo, Norway

	Boys					Girls				
	<i>n</i>	Suicidal ideation (%)	95% CI	<i>p</i> -value	Missing <i>n</i>	<i>n</i>	Suicidal ideation (%)	95% CI	<i>p</i> -value	Missing <i>n</i>
All	1,621	7.4	6.1–8.7			2,061	13.7	12.2–15.2		
Itch										
No	1,190	5.9	4.6–7.2	<0.001	32	1,327	10.7	9.0–12.4	<0.001	59
A little	302	8.6	5.4–11.8			456	17.3	13.8–20.8		
Severe	97	20.6	12.6–28.6			219	24.2	18.5–29.9		
Pain sites										
None	582	4.1	2.5–5.7	<0.001	64	263	5.3	2.6–8.0	<0.001	143
One or two	687	5.5	3.8–7.2			757	8.3	6.3–10.3		
Three to five	288	17.7	13.3–22.1			898	20.3	17.7–22.9		
Family income										
High	655	5.8	4.0–7.6	0.07	270	742	8.5	6.5–10.5	<0.001	332
Low	696	8.3	6.3–10.3			987	16.5	14.2–18.8		
Ethnicity										
Western	1,125	5.4	4.1–6.7	<0.001	149	1,428	10.6	9.0–12.2	<0.001	154
Non-Western	347	12.7	9.2–16.2			479	21.9	18.2–25.6		

*Severe itch includes self-reported quite a lot and very much itch (see Methods).

**There were five possible pain sites: head, neck/shoulder, arms/legs/shoulders, stomach, and back (see Methods).

CI: confidence interval.