

Table SI. Hyaluronan (HA) distribution in lesional and non-lesional skin of patients with psoriatic arthritis (PsA)

	All	Mono-oligo arthritis	Polyarthritis	Axial	DIP	Enthesis
HA intensity in lesional skin, <i>n</i>						
Epidermis > dermis	33	7	17	3	1	5
Epidermis = dermis	3	1	2	0	0	0
Epidermis < dermis	2	0	2	0	0	0
Unable to decide	13	3	3	1	0	6
HA localization in basement membrane zone of lesional skin, <i>n</i>						
Devoid of HA	22	3	13	3	1	2
Partly devoid of HA	9	3	3	0	0	3
Normal	1	0	1	0	0	0
Unable to decide	19	5	7	1	0	6
HA intensity in non-lesional skin, <i>n</i>						
Epidermis > dermis	11	1	7	1	0	2
Epidermis = dermis	13	6	5	1	0	1
Epidermis < dermis	15	2	8	1	1	3
Unable to decide	12	2	4	1	0	5
HA localization in basement membrane zone of non-lesional skin, <i>n</i>						
Devoid of HA	11	2	6	0	0	3
Partly devoid of HA	22	6	11	1	1	3
Normal	5	1	2	2	0	0
Unable to decide	12	2	5	1	0	4

DIP: distal joint involvement.