

Table SII. Other topical therapies included in the meta-analysis

| Study | Intervention | Type of treatment | Treatment duration (days) | Sample size in active treatment arm <i>n</i> | Sample size in control treatment arm <i>n</i> | Trial design |
|------------------------------------|--|-------------------------|---------------------------|--|---|--------------|
| Gueniche et al., 2008 (51) | V. filiformis cream 5% vs. vehicle | Non-pathogenic bacteria | 30 | 37 | 38 | Parallel |
| Lee et al., 2008 (16) | Rosmarinic acid 0.3% cream vs. vehicle | Anti-inflammatory | 56 | 21 | 21 | Left/right |
| Griffiths et al., 2002 (6) | Cipamfylline 0.15% cream vs. vehicle | Anti-inflammatory | 14 | 54 | 54 | Left/right |
| Patzelt-Wenzler et al., 2000 (44) | Kamillosan cream vs. vehicle | Anti-inflammatory | 14 | 69 | 33 | Left/right |
| Mayser et al., 2006 (45) | Ciclopiroxolamine 1% cream vs. vehicle | Anti-fungal | 28 | 16 | 13 | Parallel |
| Patrizi et al., 2008 (46) | MAS063DP vs. vehicle | Non-steroidal cream | 43 | 20 | 20 | Parallel |
| Abramovits et al., 2006 (47) | MAS063DP vs. vehicle | Non-steroidal cream | 50 | 145 | 73 | Parallel |
| Boguniewicz et al., 2008 (48) | MAS063DP vs. vehicle | Non-steroidal cream | 43 | 72 | 70 | Parallel |
| Giordano-Labadie et al., 2006 (49) | Exomega milk + cleansing bar vs. cleansing bar alone | Moisturizer | 56 | 37 | 39 | Parallel |
| Klövekorn et al., 2007 (50) | Ointment containing extracts from Mahonia aquifolium, Viola tricolor, and Centella asiatica (5 g/100 g ointment) vs. vehicle | Herbal | 28 | 88 | 88 | Left/right |