

Table SI. Characteristics of patients with *Helicobacter cinaedi* bacteraemia accompanied by skin lesions

| Case No. | Age | Sex | Underlying disease | Blood culture (<i>H. cinaedi</i>) | Fever | Diarrhoea | Location of skin lesions | Erythemas n | Size range of erythema cm | Eruption | Pain | Treatment | Recurrence |
|----------|-----|-----|---|-------------------------------------|-------|-----------|--------------------------|-------------|---------------------------|---|------|---|------------|
| 1 | 63 | M | Chronic renal failure | Positive | + | – | Arms, legs | 5 | 3–8 | Dark-red erythemas | + | <i>Clarithromycin</i> , cefdinir, amoxicillin | None |
| 2 | 79 | M | Lung cancer | Positive | + | + | Arms, legs | 12 | 3–7 | Violaceous erythemas | – | Amoxicillin, meropenem | None |
| 3 | 62 | M | Lung cancer | Positive | + | – | Legs | 3 | 6–10 | Infiltrated erythematous plaques | + | Ampicillin/sulbactam | None |
| 4 | 56 | F | Lymphoma | Positive | + | – | Legs | 2 | 5–7 | Reddish-to-violaceous erythema | + | Amoxicillin, cefepime, ampicillin/sulbactam | >4 times |
| 5 | 68 | F | Chronic renal failure, Sjögren syndrome | Positive | + | – | Arms, legs | 4 | 5–10 | Diffuse dark erythema | + | Cefazidime, fosfomycin | 1 time |
| 6 | 37 | F | Lymphoma | Positive | + | + | Left leg | 2 | 5–7 | Erythema | + | Amoxicillin, cefepime | 2 times |
| 7 | 71 | M | Autoimmune haemolytic anaemia | Positive | + | – | Thighs | 2 | 12–14 | Erythema | + | Amoxicillin, ampicillin/sulbactam | 3 times |
| 8 | 57 | F | Lymphoma | Positive | + | – | Left arm, left leg | 2 | ND | Pink-red infiltrated erythematous plaques | + | Cephazolin, panipenem | None |
| 9 | 64 | F | Lymphoma | Positive | + | – | Legs | 6 | 2–3 | Pink-red infiltrated erythema | + | Amoxicillin, levofloxacin | 3 times |

ND: not described.