

Table SV. Development and association between nodules and pruritus. *p*-values were obtained from McNemar's exact test (intra-group comparisons) and from Fisher's exact test (inter-group comparisons)

Groups	Presence of nodules				Presence of pruritus				<i>p</i> -value					
	After one year		<i>p</i> -value		Baseline		After one year		Intra ^a		Inter ^b			
	Baseline	year	Intra		SA	Inv	SA	Inv	SA	Inv	SA	Inv	SA	Inv
Exposed	0/130	23/130	<0.001 (0/130 vs. 23/130)		12/94	2/98	56/94	10/98	<0.001 (12/94 vs. 56/94)	0.039 (2/98 vs. 10/98)				
Unexposed	0/72	NA			4/50	0/68	NA	NA						
Exposed vs. unexposed:														
(i) Baseline vs. baseline				>0.3									>0.3	>0.3
(ii) Baseline vs. 1 year				(0/72 vs. 0/130)									(4/50 vs. 12/94)	(0/68 vs. 2/98)
				<0.001									<0.001	0.006
				(0/72 vs. 23/130)									(4/50 vs. 56/94)	(0/68 vs. 10/98)

^aComparisons baseline vs. after one year within the group of the exposed. ^bComparisons between exposed and unexposed. SA: Self-assessed; Inv: Investigator-assessed.