

Table SI. Socio-demographic characteristics in patients with mycosis fungoides (MF) and controls (Clalit Health Services database: population-based cohort, Rabin Medical Center: institution-based cohort)

	Population-based cohort			Institution-based cohort		
	Patients n = 683 n (%)	Controls n = 1,700 n (%)	p-value	Patients n = 343 n (%)	Controls n = 846 n (%)	p-value
Age group						
5–14 years	5 (0.7)	14 (0.8)	0.969	7 (2.0)	16 (1.9)	0.361
15–24 years	11 (1.6)	32 (1.9)		8 (2.3)	28 (3.3)	
25–34 years	28 (4.1)	72 (4.2)		22 (6.4)	66 (7.8)	
35–44 years	27 (4.2)	74 (4.4)		23 (6.7)	74 (8.8)	
45–54 years	65 (10.1)	176 (10.4)		42 (12.2)	114 (13.5)	
55–64 years	132 (20.0)	344 (20.2)		79 (23.0)	186 (22)	
65–74 years	182 (26.9)	458 (26.9)		82 (23.9)	178 (21.0)	
75–84 years	182 (24.9)	412 (24.2)		58 (16.9)	134 (15.8)	
≥85 years	51 (7.1)	118 (6.9)		22 (6.4)	50 (5.9)	
Age range (mean ± SD)	7–98 (66.05 ± 16.14)	7–110 (65.2 ± 16.0)		7–97 (60.92 ± 18.14)	10–99 (59.86 ± 18.20)	
Gender						
Male	438 (64.1)	1,088 (64.0)	0.496	229 (66.8)	572 (67.6)	0.414
Female	245 (35.9)	612 (36.0)		114 (33.2)	274 (32.4)	
Socio-economic status						
Low	188 (28.0)	609 (36.4)	<0.0001	72 (21.7)	222 (26.7)	0.009
Medium	333 (49.6)	645 (38.5)		129 (38.9)	359 (43.1)	
High	151 (22.5)	420 (25.1)		131 (39.5)	252 (30.3)	
Ethnicity						
Jewish	650 (95.2)	1,450 (85.3)	<0.0001	338 (98.5)	796 (94.1)	0.001
Arab	33 (4.8)	250 (14.7)		5 (1.5)	50 (5.9)	