

Table SII. Clinical and histopathological characteristics of cutaneous squamous cell carcinoma (cSCCs) with local recurrence and metastasis

	Sex	Age (years)	Immunosuppression	Treatment	Localization	Tumour diameter (mm)	Tumour thickness (mm)	Invasion deeper structures	Differentiation	Perineural invasion	Angio-invasion	Died of cSCC	Time to event (months)
<i>Local recurrence</i>													
Patient 1	F	82	No	SE	Other head/neck	9	1.8	No	Moderate	No	No	No	21
Patient 2	M	69	No	SE	Other head/neck	14	6.2	No	Good	No	No	No	35
Patient 3	F	65	Yes	SE	Upper extremities	13	1.1	No	Moderate	No	No	No	9
Patient 4	M	70	No	RT	Ear (auditory meatus)	51	7.5	No	Moderate	No	No	Yes	12
Patient 5	F	88	No	SE + RT	Other head/neck	10	5.0	No	Moderate	Yes	No	No	9
Patient 6	M	83	No	SE	Ear	30	6.0	Cartilage	Moderate	No	No	No	4
Patient 7	F	79	No	SE	Other head/neck	14	10.0	No	Poor	No	No	No	36
Patient 8	M	73	No	SE	Ear	13	2.8	No	Good	No	No	No	40
Patient 9	M	65	No	SE	Other head/neck	8	3.9	No	Good	No	No	No	53
Patient 10	M	66	No	SE + RT	Lip	15	8.5	Muscle	Moderate	Yes	No	No	3
Patient 11 ^a	F	84	No	SE	Lower extremities	29	7.5	No	Moderate	No	No	Yes	3
<i>Metastasis</i>													
Patient 1	M	80	No	SE	Ear	12	5.2	Muscle + cartilage	Poor	No	No	No	2
Patient 2	M	76	Yes	RT	Ear	21	2.2	No	Moderate	No	No	Yes	20
Patient 3	M	71	No	None	Ear	35	2.5	No	Good	No	No	Yes	0
Patient 4	M	80	No	SE	Ear	12	3.7	No	Poor	Yes	No	No	2
Patient 5	M	78	No	SE	Other head/neck	26	13.0	Muscle	Good	Yes	No	No	18
Patient 6	M	75	No	SE	Ear	23	7.3	Cartilage	Poor	No	No	No	47
Patient 7 ^a	F	84	No	SE	Lower extremities	29	7.5	No	Moderate	No	No	Yes	2

^aPatient with metastasis and local recurrence.
SE: surgical excision; RT: radiotherapy.