

Table SII. Localization of newly acquired melanocytic naevi (MN)

Localization	Intervention n (%)	Control n (%)	Total n (%)
Scalp	18 (0.8)	22 (1.0)	40 (0.9)
Face and ears	285 (12.4)	319 (14.5)	604 (13.4)
Neck, ventral	83 (3.6)	83 (3.8)	166 (3.7)
Neck, dorsal	138 (6.0)	137 (6.2)	275 (6.1)
Trunk, ventral	175 (7.6)	155 (7.0)	330 (7.3)
Trunk, dorsal	315 (13.7)	294 (13.4)	609 (13.5)
Upper arm, anterior	130 (5.7)	122 (5.5)	252 (5.6)
Upper arm, posterior	225 (9.8)	199 (9.0)	424 (9.4)
Lower arm, anterior	101 (4.4)	83 (3.8)	184 (4.1)
Lower arm with elbow, posterior	249 (10.8)	247 (11.2)	496 (11.0)
Thigh, anterior	167 (7.3)	154 (7.0)	321 (7.1)
Thigh, posterior	69 (3.0)	48 (2.2)	117 (2.6)
Lower legs with knee, anterior	126 (5.5)	120 (5.4)	246 (5.5)
Lower legs, posterior	71 (3.1)	83 (3.8)	154 (3.4)
Palmar	11 (0.5)	7 (0.3)	18 (0.4)
Back of the hands	101 (4.4)	88 (4.0)	189 (4.2)
Dorsum of the feet	27 (1.2)	34 (1.5)	61 (1.4)
Plantar	5 (0.2)	6 (0.3)	11 (0.2)
Total	2,296 (51.0)	2,202 (49.0)	4,498

$p=0.66$ (χ^2 test).