

Table SII. Comparative effectiveness of patients with and without psychiatric disease (interdisciplinary treatment group)

	Interdisciplinary treatment group (n=23)	
	With psychiatric disorder (n=10)	Without psychiatric disorder (n=13)
Primary outcome		
Quality of life response		
Change in DLQI, mean \pm SD	14.8 \pm 9.7 [n=9]	7.3 \pm 9.1 [n=12]
Secondary outcome		
Quality of life response		
Proportion with reduction in DLQI \geq 5 units ^a , n (%)	8/9 (88.9) [n=9]	6/12 (50.0) [n=12]
Clinical response		
Change in PASI, mean \pm SD	3.7 (6.8)	9.7 (7.2)
PASI75 response, n (%)	2/10 (20.0)	6/13 (46.2)
PASI50 response, n (%)	4/10 (40.0)	8/13 (61.5)
Global disease severity		
IGA clear/almost clear, n (%)	4/9 (44.4)	6/11 (54.6)
Patient satisfaction ^b		
VAS-score, mean \pm SD	59.1 \pm 38.4 [n=9]	66.2 \pm 28.2 [n=13]
Change in VAS score since baseline	15.0 \pm 33.5 [n=9]	18.3 \pm 38.7 [n=13]

^aIn patients with baseline Dermatology Life Quality Index (DLQI) \geq 5. ^bAssessed using a 100 unit visual analogue scale (VAS) with 0 reflecting total dissatisfaction and 100 reflecting maximal satisfaction with treatment/medical care of atopic eczema. None of the observed differences between groups was statistically significant.