

Table SII. Demographic and clinico-pathological findings by site-specific melanoma. Crude odds ratio (OR) of developing melanoma skin cancer on the trunk and extremity (n = 108)

	Extremity (n=51) n (%)	Trunk (n=57) n (%)	OR (95% CI)
Breslow depth			
≤0.90 mm	47 (92.2)	49 (86.0)	–
≥0.91 mm	4 (7.8)	8 (14.0)	1.92 (0.53–6.88)
Clark level			
1	4 (7.8)	3 (5.3)	0.99 (0.15–6.04)
2	29 (56.9)	22 (38.6)	–
3–4	18 (35.3)	32 (56.1)	2.34 (0.98–5.66)
Not neoplastic dermatoses			
No	43 (84.3)	53 (93.0)	–
Yes	8 (15.7)	4 (7.0)	0.41 (0.09–1.63)
Phototype			
III+IV	27 (52.9)	35 (61.4)	–
II+I	24 (47.1)	22 (38.6)	0.7 (0.33–1.52)
Hair colour			
Black, dark brown	24 (47.1)	31 (54.4)	–
Brown	19 (37.2)	17 (29.8)	0.69 (0.27–1.75)
Blond/red	8 (15.7)	9 (15.8)	0.87 (0.26–2.96)
Eye colour			
Dark brown, brown	36 (70.6)	43 (75.4)	–
Green/ Blue	15 (29.4)	14 (24.6)	0.78 (0.31–1.99)
Total melanocytic naevus count			
< 10	20 (39.2)	27 (47.4)	–
10–50	14 (27.5)	16 (28.1)	0.85 (0.30–2.35)
> 50	17 (33.3)	14 (24.5)	0.61 (0.22–1.68)

Risks of malignant melanoma localised in the trunk. CI: confidence intervals.