

Table SI. Main characteristics of the 12 cases of acral localised cutis laxa

References	Sex/Age		Urticaria	Joint hyperlaxity	Monoclonal gammopathy	Amyloid deposits in	
	(years)	Localisation				skin lesion	Hand X-ray
Fischer et al. (1)	F/27	Hands, feet	+	-	-	-	Normal
Yoneda et al. (2)	M/62	Hands, feet	-	-	IgG κ, myeloma	+	Osteoporosis
Yoneda et al. (2)	M/71	Hands	-	-	IgG κ, myeloma	+	Not available
Klein-Lerousseau et al. (3)	F/60	Hands, feet	-	-	Not available	-	Not available
Klein-Lerousseau et al. (3)	M/60	Hands	-	-	Not available	Not available	Not available
Martin et al. (4)	M/55	Hands, feet	+	-	-	-	Normal
Rongioletti et al. (5)	M/62	Hands, feet	+	-	-	-	Erosive subarticular cystic changes (related to rheumatoid arthritis)
Appiah et al. (6)	F/64	Hands	-	-	κ light chain, myeloma	+	Not available
Ferrandiz-Pulido et al. (7)	M/63	Hands	-	-	λ light chain, myeloma	+	Not available
Dicker et al. (8)	F/59	Hands	-	-	Not typed, myeloma	+	Normal
Marti et al. (9)	F/55	Hands, feet	-	-	-	-	Normal
Present case	M/36	Hands, feet	+	+	IgA λ, myeloma	-	Dislocation