

Table SII. Comparison of reflectance confocal microscopy (RCM) findings during the course of treatment. Only positive findings are shown

RCM finding <i>n</i> = 14	Time frame difference (mean score for the addition of 4 areas)	
		<i>p</i> -value
Scaling	t1 (5.07)–t2 (2.14)	0.001
	t3 (1.64)–t4 (1.29)	0.535
	t1 (5.07)–t3 (1.64)	0.001
	t1 (5.07)–t4 (1.29)	0.000
	t2 (2.14)–t3 (1.64)	0.303
Polygonal nucleated cells at stratum corneum	t1 (5.28)–t2 (7.21)	0.003
	t3 (7.14)–t4 (5.50)	0.014
	t1 (5.28)–t3 (1.64)	0.001
	t1 (5.28)–t4 (5.50)	0.696
	t2 (7.21)–t3 (1.64)	0.900
Detached corneocytes	t1 (2.86)–t2 (5.36)	0.004
	t3 (4.65)–t4 (3.79)	0.118
	t1 (2.86)–t3 (4.65)	0.050
	t1 (2.86)–t4 (3.79)	0.311
	t2 (5.36)–t3 (4.65)	0.273
Area of atypical honeycomb	t1 (11.79)–t2 (8.93)	0.002
	t3 (7.78)–t4 (7.71)	0.927
	t1 (11.79)–t3 (7.78)	0.000
	t1 (11.79)–t4 (7.71)	0.000
	t2 (8.93)–t3 (7.78)	0.063
Degree of atypia in typical honeycomb	t1 (7.07)–t2 (4.93)	0.000
	t3 (4.71)–t4 (4.14)	0.311
	t1 (7.07)–t3 (4.71)	0.002
	t1 (7.07)–t4 (4.14)	0.000
	t2 (4.93)–t3 (4.71)	0.657
Inflammation	t1 (3.43)–t2 (5.00)	0.002
	t3 (5.93)–t4 (5.21)	0.441
	t1 (3.43)–t3 (5.93)	0.004
	t1 (3.43)–t4 (5.21)	0.025
	t2 (5.00)–t3 (5.93)	0.187