

Table SIII. Patients' demographic and clinical characteristics (n = 105). Percentages may not add up to 100% due to missing responses and/or the possibility of giving multiple answers

	n (%)
Gender	
Male	44 (42)
Female	61 (58)
Age, mean ± SD	42.4 (13.2)
Age, median (inter quartile range (IQR))	42 (22.5)
Marital status	
Unmarried	31 (29.2)
Married or cohabiting	63 (59.4)
Widowed	1 (0.9)
Divorced	10 (9.5)
Education	
No education	–
Primary school	3 (2.8)
Lower general secondary education	20 (18.9)
Intermediate vocational education	46 (43.4)
Secondary education	6 (5.7)
Higher vocational education	19 (18.1)
University	11 (10.4)
Ethnicity	
Caucasian	88 (83.4)
Surinam/Hinduistic	5 (4.8)
Turkish	3 (2.9)
Moroccan	–
Antillean	–
African	–
Chinese	1 (1.0)
Other	7 (6.7)
Work	
Yes	72 (67.9)
No, reason:	33 (31.4)
Illness	4 (3.8)
Unemployed	1 (0.9)
Disabled	13 (12.3)
Retired	7 (6.5)
Student	6 (5.7)
Other	4 (4.7)
Diagnosis	
Psoriasis	52 (49.5)
Atopic dermatitis	25 (23.8)
Acne	2 (1.9)
Vitiligo	6 (5.7)
Hidradenitis suppurativa	14 (13.3)
Other	8 (7.6)
Disease duration (n=100), months, mean ± SD	19.7 ± 64.4
Disease duration, months, median (IQR)	1.71 (3)
Disease severity	
Patient <sup>a</sup>	
Mean ± SD	3.1 ± 1.03
Median (IQR)	3 (2)
Health care provider <sup>b</sup>	
Mean ± SD	3.13 ± 0.90
Median (IQR)	3 (1)
Global impairment of HRQoL	
Patient <sup>c</sup>	
Mean ± SD	3.3 ± 1.1
Median (IQR)	3 (1)
Health care provider <sup>d</sup>	
Mean ± SD	2.7 ± 0.8
Median (IQR)	3 (1)
Reason for inclusion	
Negative impact of skin disease on HRQoL	74 (70.5)
High motivation to learn about HRQoL	45 (42.9)
Disease severity	40 (38.1)
Other	5 (4.8)

Global disease severity from <sup>a</sup>the patients' (n = 104) and <sup>b</sup>the health care providers' perspective (5-point scale, 1: not severe to 5: very severe); <sup>c</sup>Patient (n = 103) and <sup>d</sup>health care provider (n = 104) global assessment of impairment of patients' health related quality of life (HRQoL) (1: no impairment to 5: severe impairment).