

Table SII. Distribution characteristics of lichenification by study, by treatment and by week. For a description of the studies see Material and Methods

Week	Study	Treatment	n	Mean ± SD	Median	Min–Max
0	FLT001	OD	137	1.93 ± 0.65	2	0–3
		BD	133	1.92 ± 0.60	2	0–3
		Total	270	1.92 ± 0.63	2	0–3
	FLT002	OD	123	2.22 ± 0.52	2	1–3
		BD	122	2.24 ± 0.49	2	1–3
		Total	245	2.23 ± 0.50	2	1–3
	FLT401	Placebo	76	1.66 ± 0.89	2	0–3
		OD	79	1.63 ± 0.82	2	0–3
		BD	79	1.68 ± 0.83	2	0–3
		Total	234	1.66 ± 0.85	2	0–3
	Total	Placebo	76	1.66 ± 0.89	2	0–3
		OD	339	1.96 ± 0.69	2	0–3
		BD	334	1.98 ± 0.66	2	0–3
		Total	749	1.94 ± 0.71	2	0–3
1	FLT001	OD	132	1.38 ± 0.71	1.5	0–3
		BD	125	1.36 ± 0.71	1.5	0–3
	FLT002	OD	118	1.64 ± 0.68	1.5	0–3
		BD	118	1.57 ± 0.71	1.5	0–3
	FLT401	Placebo	74	1.39 ± 0.79	1.5	0–3
		OD	77	1.01 ± 0.68	1	0–3
		BD	77	0.91 ± 0.74	1	0–3
	Total	Placebo	74	1.39 ± 0.79	1.5	0–3
		OD	327	1.38 ± 0.73	1.5	0–3
		BD	320	1.33 ± 0.76	1.5	0–3
2	FLT001	OD	122	0.96 ± 0.74	1	0–2.5
		BD	117	0.92 ± 0.70	1	0–3
	FLT002	OD	104	1.26 ± 0.74	1	0–3
		BD	107	1.22 ± 0.69	1	0–3
	FLT401	Placebo	63	1.13 ± 0.92	1	0–3
		OD	73	0.70 ± 0.65	0.5	0–2.5
		BD	74	0.68 ± 0.64	0.5	0–2.5
	Total	Placebo	63	1.13 ± 0.92	1	0–3
OD		299	1.00 ± 0.75	1	0–3	
BD		298	0.97 ± 0.71	1	0–3	
3	FLT001	OD	109	0.78 ± 0.69	1	0–2.5
		BD	111	0.76 ± 0.67	0.5	0–3
	FLT002	OD	96	1.07 ± 0.72	1	0–3
		BD	92	1.13 ± 0.73	1	0–3
	FLT401	Placebo	56	1.01 ± 0.88	1	0–3
		OD	69	0.54 ± 0.58	0.5	0–2.5
		BD	69	0.56 ± 0.63	0.5	0–2.5
	Total	Placebo	56	1.01 ± 0.88	1	0–3
OD		274	0.82 ± 0.70	1	0–3	
BD		272	0.83 ± 0.71	1	0–3	
4	FLT001	OD	99	0.65 ± 0.64	0.5	0–2
		BD	98	0.59 ± 0.60	0.5	0–2
	FLT002	OD	83	1.02 ± 0.64	1	0–2.5
		BD	80	1.07 ± 0.71	1	0–2.5
	FLT401	Placebo	53	0.95 ± 0.86	1	0–3
		OD	62	0.38 ± 0.54	0	0–2.5
		BD	57	0.40 ± 0.54	0	0–2.5
	Total	Placebo	53	0.95 ± 0.86	1	0–3
OD		244	0.71 ± 0.67	0.5	0–2.5	
BD		235	0.71 ± 0.68	0.5	0–2.5	

OD: once daily; BD: twice daily; mean: mean lichenification score; SD: standard deviation; Min: minimum lichenification score; Max: maximum lichenification score.

Lichenification criteria: 0: absent, 0.5: very mild, 1.0: mild, 1.5: mild to moderate, 2.0: moderate, 2.5: severe, 3.0: very severe.