

Table SI. Patient clinical data

ID	Sex/Age	Psoriasis since	Biopsy site	Localisation of lesions	Skin type	Given dose	Treatments <i>n</i>	Response
1	M/40	>20 years	Buttock, lower back	Whole body	I–II	0.3–1.4 J	34	Good
2	F/73	Unknown	Arms	Arms, back, soles, scalp	III	0.3–2.0 J	20	Excellent
3	M/59	>20 years	Thighs	Legs, buttocks, left elbow, hands	III	0.3–1.7 J	23	Excellent
4	M/67	<1 years	Buttocks, thigh	Elbows, legs, scalp, buttocks	II–III	0.3–1.5 J	20	Excellent
5	M/26	1 year	Buttocks	Legs, arms, scalp, face, buttocks	I–II	0.1–0.55 J	15 <sup>a</sup>	Unknown
6	M/17	>10 years	Buttocks	Whole body, scalp	IV	0.3–1.7 J <sup>b</sup>	21	Unsatisfactory
7	F/47	>10 years	Buttocks	Whole body	I	0.3–2.2 J	23	Unsatisfactory
8	M/43	>10 years	Buttocks	Lower legs, arms, trunk	III	0.3–2.6 J	25	Excellent
9	F/46	>10 years	Legs	Legs, elbows, hands	III	0.3–2.6 J	30	Good
10	F/74	2 years	Buttocks	Whole body	II	0.3–2.1 J	20	Excellent
11	F/33	>10 years	Buttocks	Whole body	III	0.3–2.0 J	21	Excellent
12	F/28	>10 years	Buttocks	Whole body	I	0.2–1.0 J	19	Excellent

<sup>a</sup>Stopped treatment half way. <sup>b</sup>0.3–1.7 J, stop 3 weeks, 0.9–1.3 J.