

Table SII. History of human papillomavirus (HPV)-related diseases and warts after 24 years

Variable	Physicians (n=158) n (%)	Non-physicians (n=129) n (%)	p	CO ₂ laser+ (n=166) n (%)	CO ₂ laser- (n=121) n (%)	p	Dermato-venerology (n=100) n (%)	Gynaecology (n=187) n (%)	p
History of HPV-related diseases	96 (60.8)	87 (67.4)	0.27	102 (61.4)	81 (66.9)	0.38	73 (73.0)	110 (58.8)	0.02
Warts on hands	39 (24.7)	31 (24.0)	1.00	40 (24.1)	30 (24.8)	1.00	31 (31.0)	39 (20.9)	0.06
Warts on feet	62 (39.2)	46 (35.7)	0.54	61 (36.7)	47 (38.8)	0.81	44 (44.0)	64 (34.2)	0.12
Genital warts	16 (10.1)	16 (12.4)	0.58	18 (10.8)	14 (11.6)	0.85	11 (11.0)	21 (11.2)	1.00
Nasal or oro-pharyngeal warts	2 (1.3)	1 (0.8)	1.00	2 (1.2)	1 (0.8)	1.00	3 (3.0)	0 (0.0)	0.04
Cervical dysplasia/cancer ^a	11 (11.4)	17 (13.3)	0.84	12 (9.9)	16 (15.5)	0.23	7 (9.6)	21 (13.9)	0.52
Warts after 24 years of age	36 (22.7)	32 (24.8)	0.78	39 (21.7)	29 (24.0)	1.00	27 (27.0)	41 (21.9)	0.38
Warts on hands	20 (12.7)	15 (11.6)	0.86	22 (13.3)	13 (10.7)	0.59	18 (18.0)	17 (9.1)	0.04
Warts on feet	24 (15.2)	17 (13.2)	0.62	22 (13.3)	19 (15.7)	0.69	13 (13.0)	28 (15.0)	0.73
Genital warts	6 (3.8)	3 (2.3)	0.52	4 (2.4)	5 (4.1)	0.50	5 (5.0)	4 (2.1)	0.28
Nasal or oral	2 (1.3)	1 (0.8)	1.00	2 (1.2)	1 (0.8)	1.00	3 (3.0)	0 (0.0)	0.04

^a% of woman.

p; Fisher's exact test.