Supplementary material to article by K. Kofoed et al. "Low Prevalence of Oral and Nasal human papillomavirus in Employees Performing CO<sub>2</sub>-laser Evaporation of Genital Warts or Loop Electrode Excision Procedure of Cervical Dysplasia"

Table SI. Baseline characteristics of the included healthcare personnel

Characteristic	Value n (%)
Sex, n (%)	(, 0)
Male	64 (22.3)
Female	223 (77.7)
Age, median years, range	44 (22–68)
Occupation, n (%)	(== 00)
Physician	158 (55.1)
Nurse	122 (42.5)
Other <sup>a</sup>	7 (2.4)
Work place, <i>n</i> (%)	
Department of Dermato-venerology	100 (34.8)
Department of Gynaecology	187 (65.2)
Participated in $CO_2$ laser treatment of genital warts, $n$ (%)	166 (57.8)
median years, range	5 (1–25)
Performed cryotherapy of genital warts, $n$ (%)	95 (33.1)
median years, range	5 (1–25)
Participated in electrosurgical treatment of genital warts, $n$ (%)	104 (36.2)
median years, range	5 (1–35)
Participated in loop electrosurgical excision procedure, $n$ (%)	112 (39.0)
median years, range	7 (1–25)
Vaccinated with quadrivalent human papillomavirus vaccine, $n$ (%)	9 (3.1)

<sup>&</sup>lt;sup>a</sup>researcher, lab technician, nursing assistant, medical and nurse student.