

Table SI. Demographic characteristics and clinical assessment of patients with acquired cold urticaria

Patient (sex/age, years)	Disease duration, years	Critical stimulation time threshold (s)			Critical temperature threshold (°C)			Evolution
		Pre-	Post-	Elongation n (%)	Pre-	Post-	Reduction n (%)	
1 (F/48)	1	90	180	90 (50)	14	10	4 (29)	Active
2 (F/38)	12	210	>300	91 (30)	8	<4	5 (63)	Cure
3 (F/37)	2.5	150	>300	151 (50)	14	6	7 (57)	Active
4 (F/55)	22	210	>300	91 (30)	12	<4	9 (75)	Cure
5 (F/52)	2.5	150	>300	151 (50)	18	<4	15 (83)	Cure
6 (M/51)	8.5	210	>300	91 (30)	4	<4	1 (25)	Cure
7 (M/43)	3.5	150	>300	151 (50)	16	<4	13 (81)	Cure
8 (M/40)	2	<60	240	181 (60)	24	4	20 (83)	Active
9 (M/52)	1	60	180	120 (66)	20	14	6 (30)	Active
10 (F/36)	5	180	>300	121 (40)	12	<4	9 (75)	Cure
11 (M/62)	1.5	60	>300	241 (80)	10	<4	7 (70)	Cure
12 (F/40)	2	300	>300	1 (0)	4	<4	1 (25)	Cure
13 (M/53)	3	180	210	30 (14)	14	10	4 (29)	Active
14 (M/40)	10	210	>300	81 (27)	20	<4	17 (85)	Cure
15 (F/34)	2	60	60	0 (0)	26	24	2 (7)	Active
16 (F/28)	8	120	>300	181 (60)	10	<4	7 (70)	Cure
17 (M/40)	1.5	180	>300	121 (40)	12	<4	9 (75)	Cure
18 (F/61)	5	240	>300	61 (20)	8	<4	7 (62)	Cure
19 (M/45)	3	120	>300	181 (60)	14	<4	11 (78)	Cure

n = difference between pre and post (% of post-values).