

Table SIV. Calculated crude odds ratios of factors and remission of atopic dermatitis (AD) based on data in Illi et al. (5)

Factors	Complete remission	Intermittent AD	Persistent AD	OR (95% CI)
Frequent scratching with early AD	12/83	26/73	26/36	0.12 (0.059–0.259)
≥ Two atopic family members	16/83	23/72	18/36	0.39 (0.200–0.762)
Parental AD	9/81	13/71	69/36	0.48 (0.209–1.115)
Parental hay fever	38/83	41/72	20/3	0.65 (0.366–1.157)
Parental asthma	19/83	14/72	12/36	0.81 (0.414–1.582)
Increased cord blood IgE levels ($\geq 0.9 \text{ kU/l}$)	11/80	13/68	2/34	0.92 (0.399–2.141)
Increased total IgE levels ($\geq 30 \text{ kU/l}$)	20/53	24/44	11/22	0.54 (0.257–1.120)
Any sensitization	13/52	21/42	13/22	0.31 (0.142–0.693)
Any food sensitization	7/52	16/43	8/22	0.28 (0.109–0.714)
Any inhalant sensitization	8/52	14/43	9/22	0.35 (0.141–0.861)
Early wheeze	24/77	28/66	17/32	0.53 (0.286–0.996)

Statistically significant correlations are highlighted in bold. OR: odds ratio; 95% CI: 95% confidence interval; IgE: immunoglobulin E.