

Table SII. κ coefficient of agreement for various sets of visual analogue scale (VAS) bands with respect to verbal rating scale (VRS) scores

VAS cut-off values ^a	3–6–8	3–6–9	3–7–8	3–7–9	4–6–8	4–6–9	4–7–8	4–7–9
All subjects (<i>n</i> =949)	κ (95%) r	0.56 (0.53–0.59) 0.92	0.57 (0.59–0.65) 0.92	0.57 (0.54–0.80) 0.92	0.64 (0.61–0.67) 0.60 (0.56–0.65)	0.55 (0.51–0.58) 0.91	0.61 (0.58–0.64) 0.91	0.56 (0.53–0.59) 0.9
Male (<i>n</i> =446)	κ (95%) r	0.54 (0.50–0.59) 0.91	0.59 (0.54–0.63) 0.91	0.55 (0.51–0.60) 0.9	0.60 (0.56–0.65) 0.66 (0.62–0.70)	0.53 (0.49–0.58) 0.89	0.58 (0.53–0.63) 0.89	0.60 (0.55–0.64) 0.66 (0.61–0.70)
Female (<i>n</i> =503)	κ (95%) r	0.56 (0.52–0.61) 0.93	0.65 (0.61–0.69) 0.93	0.58 (0.53–0.62) 0.93	0.66 (0.62–0.70) 0.66 (0.62–0.70)	0.56 (0.51–0.90) 0.92	0.64 (0.60–0.68) 0.92	0.57 (0.52–0.61) 0.92

^aA VAS cut-off scheme of (3–6–8) refers to the following set of cut-off values: 0<3: mild pruritus; ≥3 to <6: moderate pruritus; ≥6 to <8: severe pruritus; ≥8: very severe pruritus. Bold indicates bands with the best agreement.