

Table SI. Clinical data for patients with ichthyosiform mycosis fungoides (IMF)

Patient	Age, years/ Patient Sex	Duration, years, Medical history before MF	Symptom	Clinical diagnosis	Age at onset of MF, years	eruptions, years	ichthyosiform lesions	Location of additional MF lesions	Associated lymphoma	TNM/stage
1	15/M	2	Pruritic (VAS=3)	Psoriasis, Parapsoriasis, MF	13	13	Trunk, buttocks, upper and lower limbs	Patch	–	T ₂ N ₀ M ₀ /IB
2	8/F	5	Pruritic (VAS=4)	Ichthyosis vulgaris	3	3	Scalp, face, trunk, buttocks, upper and lower limbs	None	–	T ₂ N ₀ M ₀ /IB
3	17/F	10	Asymptomatic	Ichthyosis vulgaris	7	7	Trunk, buttocks, upper and lower limbs	None	–	T ₂ N ₀ M ₀ /IB
4	63/M	DM, Fatty liver	2	Pruritic (VAS=8)	Ichthyosis vulgaris, extoliative dermatitis	61	61	Trunk, buttocks, upper and lower limbs	Plaques	–
5	30/M	BPH	5	Asymptomatic	MF	25	25	Trunk, buttocks, upper and lower limbs	Multiple ulcerated lesions	–
6	47/M	Tenosynovitis	3	Pruritic (VAS=6)	MF	44	49	Trunk, upper and lower limbs	Patches and plaques	T ₂ N ₁ M ₀ /IIA
7	28/M	Pulmonary Tuberculosis	10	Asymptomatic	Ichthyosiform MF, Hypopigmented MF	18	18	Trunk, lower limbs	Hypopigmented lesions	T ₂ N ₀ M ₀ /IB
8	29/M		3	Pruritic (VAS=7)	MF	26	30	Trunk, buttocks, upper and lower limbs	Patch and follicular lesions	PCALCL
9	26/F		6	Asymptomatic	Ichthyosiform MF, Ichthyosis vulgaris	20	20	Trunk, lower limbs	None	T ₂ N ₀ M ₀ /IB
10	57/M		20	Asymptomatic	Ichthyosis vulgaris, Ichthyosiform MF	37	37	Face, trunk, buttocks, upper and lower limbs	None	T ₂ N ₀ M ₀ /IB

DM: diabetes mellitus; BPH: benign prostatic hyperplasia; MF: mycosis fungoides; PCALCL: primary cutaneous anaplastic large cell lymphoma; M: male; F: female; VAS: visual analogue scale; TNM: tumor-node-metastasis.