

Table SII. Patient characteristics

Pat. No.	Sex/age, years	Location	Size and number of tumours	T stage	Radiotherapy dose (cGy)	Response
1	F/48	Right periumbilical abdomen	Solitary tumour 1 cm	1a	3,600	Complete remission
2	M/45	Bilateral cheek	Grouped of nodules occupying up to 4 cm and 5 cm at right and left cheek, respectively	2a	4,000	Complete remission
3	M/37	Left thigh	Solitary tumour 4 cm	1a	4,000	Complete remission
4	M/40	Left hand	Solitary tumour 2 cm	1a	3,600	Complete remission
5	M/51	Left knee and thigh	Solitary tumour 3.5 cm	1a	5,040	Partial remission
6	M/46	Right thigh	2 tumours of 2 cm and 3 cm	2a	3,600	Complete remission
7	M/16	Left buttock	Solitary tumour 6 cm	1b	4,000	Complete remission
8	F/29	Left thigh	Solitary tumour 3 cm	1a	5,000	Complete remission
9	M/22	Right thigh	Grouped of nodules occupying up to 7 cm	2a	4,400	Complete remission

T stage: solitary skin involvement <5 cm (1a), >5 cm (1b); 2a: regional skin involvement: multiple lesions limited to 1 body region or 2 contiguous body regions with all-disease-encompassing in a <15-cm diameter circular area. Patients were staged according to the tumour, node, metastases classification system according to proposals of the International Society of Cutaneous Lymphomas and European Organization for Research and Treatment of Cancer.