

Table SIII. Characteristics of the study cohort

Category	n (%)
Age, years	
Mean (standard deviation)	69.2 (12.0)
Median (min–max; interquartile range)	71 (37–103; 62–77)
<60 years	28 (22.6)
60–69 years	25 (20.2)
70–79 years	49 (39.5)
>80 years	21 (16.9)
Sex	
Female	54 (43.5)
Male	70 (56.5)
Marital status	
Living with a partner	87 (70.2)
Living alone	37 (29.8)
Professional qualification	
No vocational degree	21 (16.9)
Completed vocational education	90 (72.6)
(Technical) college degree	13 (10.5)
Working status	
Working	24 (19.4)
Not working	100 (80.6)
Monthly household income	
<1,500 €	32 (25.8)
1,500–3,000 €	53 (42.7)
>3,000 €	17 (13.9)
Not determined	22 (17.7)
Number of tumours	182
Mean (standard deviation)	1.47 (1)
Median (min–max; interquartile range)	1 (1–6; 1–2)
1	91 (73.4)
2	20 (16.1)
3	8 (6.5)
4	0 (0)
5	3 (2.4)
6	2 (1.6)
Tumour localization	
Head or neck	97 (78.2)
Body	14 (11.3)
Both head or neck and body	11 (8.9)
Not determined	2 (1.6)
Tumour size ^a	
<1 cm	72 (58.1)
1–2 cm	42 (33.9)
2.1–5 cm	8 (6.5)
>5 cm	2 (1.6)
Histological subtype	
Superficial BCC	19 (11.4)
Nodular BCC	78 (42.9)
Infiltrating BCC	58 (31.9)
Sclerodermiform BCC	4 (2.2)
Not determined	4 (2.2)
Second tumour other than BCC ^b	19 (10.4)
Recurrence	
Yes	18 (14.1)
No	105 (84.7)
Unknown	1 (0.8)
History of skin cancer	
Yes	53 (42.7)
1 previous skin cancer	18 (14.5)
>1 previous skin cancer	32 (25.8)
No	71 (57.3)

^aThe size of the largest tumour was documented in each participant. ^bFive participants were diagnosed with a squamous cell carcinoma (SCC), 2 with a collision of basal cell carcinoma (BCC) and SCC, 2 with Bowen's disease 4 with actinic keratosis, 4 with solar elastosis, one with seborrheic keratosis and one with a histiocytoma as second tumour in addition to their BCC. Percentages do not always total 100% because of rounding.