

Table SIII. Evidence table

First author (Ref), year	Design	Dosage	Outcome measures	Date of evaluation (licensed dose)	Results (mean)
Adalimumab					
Ryan (12), 2009 (n=39 ^a)	Retrospl.	Adalimumab 80 mg (W 0) followed by 40 mg once W (from W1) SC or adalimumab 40 mg EOW (from W1) SC. According to the response the dose interval was increased or decreased.	PASI75 Monotherapy PASI75 combination therapy	W 16 W 24 W 48 W 72 W 16 W24 W48 W72 W 12 Comparing with original PASI75: 27% PASI (licensed dose) W 24 Comparing with original PASI75: 36% PASI W 48 Comparing with original PASI75: 54% PASI W 12 Comparing with course PASI75: 13% PASI (licensed dose) W 24 Comparing with course PASI75: 7% PASI W 48 Comparing with course PASI75: 23% PASI W 16 (licensed dose) W 24 W 52 W 72	PASI75: 38% PASI75: 62% PASI75: 69% PASI75: 71% PASI75: 56% PASI75: 50% PASI75: 80% PASI75: 67% PASI50: 70% PASI75 PASI90 PASI (licensed dose) PASI90: 10% PASI50: 61% Comparing with original PASI75: 36% PASI90: 7% PASI50: 77% Comparing with original PASI75: 54% PASI90: 15% PASI50: 30% Comparing with course PASI75: 13% PASI90: 13% PASI50: 50% Comparing with course PASI75: 7% PASI90: 0% PASI50: 54% Comparing with course PASI75: 23% PASI90: 15% PASI50: 84% PASI75: 64% PASI90: 36% PASI50: 77% PASI75: 65% PASI90: 42% PASI50: 56% PASI75: 48% PASI90: 41% W 16 PASI50: 90% PASI75: 68% PASI90: 42% PASI50: 95% PASI75: 82% PASI90: 56% PASI75: 89% PASI90: 75% PASI50: 100% PASI75: 88% PASI90: 75% W 16 PASI75: 64% PASI90: 49% W 24 PASI75: 67% PASI90: 60% W 52 PASI75: 76% PASI90: 70% W 104 PASI75: 83% PASI90: 71% W 12 PASI50: 65% Comparing with original PASI75: 35% PASI PASI100: 0% W 24 PASI50: 69% Comparing with original PASI75: 45% PASI PASI90: 17% PASI100: 6%
van Lü mig (13), 2010 (n=30 ^a)	Prosp. ^b	Adalimumab 80 mg (W0) followed by 40 mg EOW (from W1–12) SC. From W12–48, according to the response the dose interval was maintained, increased or decreased. (Adalimumab 40 mg EOW or 40 mg every 10 days or 40 mg once W.	PASI50 PASI75 PASI90		
Warren (14), 2010 (n=46 ^a)	Retrospl.	Adalimumab 80 mg (W 0) followed by 40 mg EOW (from W1) SC. According to the response the dose was increased to 40 mg weekly the first 4 months.	PASI50 PASI75 PASI90		
Fotiadou (15), 2012 (n=52 ^a)	Retrospl.	Adalimumab 80 mg (W 0) followed by 40 mg EOW (from W1) SC. According to the response the dose interval was increased or decreased	PASI50 PASI75 PASI90		
López-Ferrer (16), 2013 (n=119 ^a)	Retrospl.	Adalimumab 80 mg (W 0) followed by 40 mg EOW (From W1) SC.	PASI75 PASI90		
van Lü mig (17), 2013 (n=85 ^a)	Prosp.	Adalimumab 80 mg (W 0) followed by 40 mg EOW (from W1) SC. According to the response the dose interval was decreased.	PASI50 PASI75 PASI90 PASI100		

			W 48	PASI50: 83%
		Comparing with original	PASI75: 57%	
		PASI	PASI90: 21%	
			PASI100: 4%	
		W 96	PASI50: 94%	
		Comparing with original	PASI75: 44%	
		PASI	PASI90: 22%	
			PASI100: 0%	
		W 132	PASI50: 83%	
		Comparing with original	PASI75: 50%	
		PASI	PASI90: 0%	
			PASI100: 0%	
		W 12	PASI50: 46%	
		Comparing with course	PASI75: 27%	
		PASI	PASI90: 8%	
			PASI100: 0%	
		W 24	PASI50: 56%	
		Comparing with course	PASI75: 31%	
		PASI	PASI90: 11%	
			PASI100: 6%	
		W 48	PASI50: 72%	
		Comparing with course	PASI75: 38%	
		PASI	PASI90: 15%	
			PASI100: 4%	
		W 96	PASI50: 72%	
		Comparing with course	PASI75: 44%	
		PASI	PASI90: 22%	
			PASI100: 0%	
		W 132	PASI50: 50%	
		Comparing with course	PASI75: 50%	
		PASI	PASI90: 0%	
			PASI100: 0%	
Menting (18), 2014 Prosp. (n=90 TE ^a)	Adalimumab 80 mg (W 0) followed by 40 mg EOW (from W1) SC or 40 mg every 10 days or 40 mg once W.	PASI75 PASI90 Non-naive	W 12 W 52 W 12 W 52	PASI75: 54% PASI90: 42.9% PASI75: 26% PASI90: 18% PASI90: 8%
<i>Etanercept</i>				
de Groot (19), 2006 Retrosp. (n=50 ^a)	Etanercept 50 mg BIW SC (W0–12).	PASI50 PASI75 PASI90 MPD MBSAD	W 12 (licensed dose)	PASI50: 71% PASI75: 22.6% PASI90: 6.5% MPD: 57.5% MBSAD: 62.3%
Berends (20), 2007 Prosp. (n=45 ^a)	Etanercept 25 mg BIW SC (W0–12). Etanercept 50 mg BIW SC (W0–12) followed by 25 mg BIW (W12–24).	PASI50 PASI75 PASI90 MPD MBSAD	W 12 W 18 W 24 W 12 W 18 W 24	PASI50: 85.7% PASI75: 21.4% PASI90: 7.1% MPD: 60% MBSAD: 54.1% PASI50: 82% PASI75: 39% PASI50: 88% PASI75: 47% PASI50: 71% PASI75: 50%
Barrera (21), 2008 Retrosp. (n=66)	Etanercept 50 mg BIW SC (W0 – 12) followed by 25 mg BIW (W12–24) or 25 mg BIW (W0–24). Thereafter one of these treatments by PASI90 losing 50% of the PASI. Three cycles of 24W. MPD	PASI50 PASI75 Cycle 1 W 12 (licensed dose)	W 12 W 18 W 24 Cycle 1 W 12 (licensed dose)	PASI50: 71% PASI75: 24% PASI50: 79% PASI75: 64% PASI50: 79% PASI75: 57% PASI50: 65.5% PASI75: 36.1% PASI90: 21.3%
			W 24	PASI50: 91.3% PASI75: 76.1% PASI90: 41.3% MPD: 78.1%

			Cycle 2 W 12	PASI50: 72.4% PASI75: 31% PASI90: 10.3%
			Cycle 2 W 24	PASI50: 94.7% PASI75: 63.2% PASI90: 31.6%
			Cycle 3 W 12	MPD: 66.9% PASI50: 60% PASI75: 33.3% PASI90: 20%
			Cycle 3 W 24	PASI50: 85.7% PASI75: 42.9% PASI90: 28.6% MPD: 76%
Driessens (22), 2008 Prospl. (n=80 ^a)	Etanercept 50 mg BIW SC (W0–12) followed by 25 mg BIW (W12–24). According to the response the dose of 25 mg BIW was temporary increased. Etanercept 25 mg BIW SC (W0–24). According to the response the dose was temporary increased.	PASI50 PASI75 PASI90 MPD PASI50 PASI75 PASI90 MPD	W 12 (licensed dose) W 24 (licensed dose) W 12 W 24	PASI50: 66% PASI75: 20% PASI90: 8% MPD: 59% PASI50: 68% PASI75: 21% PASI90: 5% MPD: 60% MPD: 74.5%
Gisondi (40), 2008 Retrospl. (n=58)	Etanercept 25 mg BIW SC.	MPD	W 24	MPD: 60%
Antoniou (23), 2009 (n=77 ^a)	Retrospl. Etanercept 50 mg BIW SC (W0–12) followed by 25 mg BIW (W12–48). According to the response the dose was increased to 50 mg BIW.	PASI50 PASI75 W 48	W 24	PASI50: 79% PASI75: 53% PASI50: 81% PASI75: 49%
Driessens (24), 2009 Prospl. (n=90 ^a)	Etanercept 50 mg BIW SC (W0–12) followed by 25 mg BIW (W12–24) or Etanercept 25 mg BIW SC (W0–W24). According to the response the dose was temporary increased to 50 mg BIW.	PASI50 PASI75 PASI90 MPD	W 24	PASI50: 69% PASI75: 39% PASI90: 18% MPD: 59.7%
Jiménez-Puya (25), 2009 (n=58)	Etanercept 50 mg BIW SC or 25 mg BIW SC.	PASI50 PASI75 PASI90 MBSAD	W 8 W 16 W 24 W 32 W 40 48W	PASI50: 77.6% PASI75: 41.4% PASI90: 6.9% PASI50: 86.8% PASI75: 66% PASI90: 37.7% PASI50: 97.5% PASI75: 85% PASI90: 57.5% PASI50: 100% PASI75: 80.8% PASI90: 57.7% PASI50: 94.7% PASI75: 84.2% PASI90: 52.6% PASI50: 100% PASI75: 92.3% PASI90: 69.2% MBSAD: 85%
Mazzotta (26), 2009 (n=234)	Prospl. ^c Etanercept 50 mg BIW SC (W0–12) followed by 25 mg BIW (W12–24)	PASI50 PASI75 MPD (psoriasis)	W 12 (licensed dose) W 24 (licensed dose)	PASI50: 77.9% PASI75: 41% MPD: 67.7% PASI50: 85.7% PASI75: 73.2% MPD: 81% PASI50: 64 % PASI75: 43% MPD: 66.7% PASI50: 79.8% PASI75: 61.8% MPD: 80.5%
Warren (27), 2009 Retrospl. (n=70 ^a)	Etanercept 25 mg BIW SC. According to the response the dose was increased to 50 mg BIW.	PASI50 PASI75 PASI90 MPD	W 12	PASI50: 67% PASI75: 35% PASI90: 7% MPD: 55%

				W 24	PASI50: 67% PASI75: 41% PASI90: 6% MPD: 59% PASI75: 57%
Antoniou (28), 2010 (<i>n</i> =35 ^a)	Retrosp. ^d	Etanercept 50 mg BIW SC (W0–12) followed by 25 mg BIW (W12–24).	PASI75	W 24 (licensed dose)	
van Lü mig (13), 2010 (<i>n</i> =30 ^a)	Prospr.	Etanercept 50 mg BIW SC or 25 mg BIW SC (W0 – 12). Thereafter dosage according to physician.	PASI50 PASI75 PASI90	W 12 W 24 W 48	PASI50: 60% PASI75: 13% PASI90: 3.3% PASI50: 58% PASI75: 19% PASI90: 3.3% PASI50: 79% PASI75: 25%
Zaragoza (29), 2010 (<i>n</i> =43)	Retrosp. ^c	Etanercept 50 mg BIW SC (W0–12) or 25 mg BIW SC (W0–12) followed by 25 mg BIW SC (W12–24).	PASI50 PASI75 MPD	W 24 W 48	PASI50: 81.4% PASI75: 60.5% MPD: 73.8% PASI50: 90.5% PASI75: 71.4% MPD: 79.6%
		According to the response the dose was increased or decreased.		W 72 W 96 W 120 W 144 W 12 W 24 W 48 W 72	MPD: 81.3% MPD: 38.2% MPD: 87.6% MPD: 88.4% PASI50: 59% PASI75: 25% PASI50: 81% PASI75: 47% PASI50: 82% PASI75: 54% PASI50: 87% PASI75: 52%
Antoniou (30), 2011 (<i>n</i> =118 ^a)	Retrosp.	Etanercept 50 mg BIW SC (W0–12) followed by 50 mg once W (from W 12).	PASI50 PASI75	W 24 W 52 W 104 W 156 W 12	PASI50: 82% PASI75: 54.1% MPD: 73.3% PASI50: 90.2% PASI75: 78.7% MPD: 85.9% PASI50: 90.2% PASI75: 83.6% MPD: 87.9% PASI50: 91.8% PASI75: 86.9% MPD: 88.4% PASI50: 91.8% PASI75: 83.6% MPD: 86.9%
Esposito (31), 2012 (<i>n</i> =61)	Retrosp. ^c	Psoriasis patients with psoriatic arthritis: Etanercept 50 mg BIW SC. Patients with only psoriasis: Etanercept 50 mg BIW, followed by 25 mg BIW SC.	PASI50 PASI75 MPD	W 24	PASI50: 82% PASI75: 54.1% MPD: 73.3% PASI50: 90.2% PASI75: 78.7% MPD: 85.9% PASI50: 90.2% PASI75: 83.6% MPD: 87.9% PASI50: 91.8% PASI75: 86.9% MPD: 88.4% PASI50: 91.8% PASI75: 83.6% MPD: 86.9%
van Lü mig (32), 2012 (<i>n</i> =152 ^a) (<i>n</i> =158 TE)	Prospr.	Etanercept 50 mg BIW SC or 25 mg BIW SC (W0–12). Thereafter dosage according to the physician.	PASI50 PASI75 PASI90	W 12 W 24 W 48 W 108 W 156 W 204 W 264 W 12 Overall	PASI50: 65.6% PASI75: 23.6% PASI90: 5.1% PASI50: 69.7% PASI75: 38.1% PASI90: 14.9% PASI75: 36.6% PASI75: 40.8% PASI75: 50% PASI75: 59.4% PASI75: 60% PASI50: 88.4% PASI75: 63.9% PASI90: 37.8% MPD: 69.6% MBSAD: 66.4% MPhGAD: 62.5% PASI75: 63% PASI90: 34.8% PASI75: 64% PASI90: 38.4%
Puig (33), 2012 (<i>n</i> =444 ^a)	Prospr.	Etanercept 50 mg BIW SC (W0 –12) or 50 mg once W or 25 mg BIW (W0–12). After 12 W patients reaching a PASI50 could continue treatment. After 6 months the decision MPD was made to treat patients continuously or intermittently.	PASI50 PASI75 PASI90 MBSAD MPhGAD	W 12 Continuous regimen W 12 Intermittent regimen	

				W 24	PASI50: 91.3% PASI75: 76.1% PASI90: 50.8% MPD: 81.4% MBSAD: 79.6% MPhGADR: 65.6%
				W 24 Continuous regimen	PASI75: 66.3% PASI90: 39.3%
				W 24 Intermittent regimen	PASI75: 79% PASI90: 54.3%
				W 36	PASI50: 73.9 PASI75: 52.6% PASI90: 32.1% MPD: 63.4% MBSAD: 66.7% MPhGADR: 46.9%
				W 36 Continuous regimen	PASI75: 68.8% PASI90: 41.9%
				W 36 Intermittent regimen	PASI75: 46.7% PASI90: 28.6% W 48
				W 48 Continuous regimen	PASI50: 83.1% PASI75: 65.6% PASI90: 39.6% MPD: 74.6% MBSAD: 75.7% MPhGADR: 59.4%
				W 48 Intermittent regimen	PASI75: 69.2% PASI90: 47.3% PhGA 0–1: 73.4% PASI75: 64.3% PASI90: 36.6% PhGA 0–1: 61.8%
Chiu (34), 2013 (n=59 ^a)	Retrosp.	Etanercept 25 mg BIW SC (W0–24).	PASI50 PASI75 PASI90 MPD	W 12	PASI50: 48% PASI75: 26% PASI90: 3% MPD: 47%
		According to the response the dose was increased to 50 mg BIW SC (for 4–12W). Thereafter dose was reduced to 25 mg BIW SC.		W 24	PASI50: 59% PASI75: 37% PASI90: 14% MPD: 61%
van Lü mig (35), 2013 (n=131 ^a) (n=134 TE)	Prospl.	Etanercept mean weekly dose 64.1 mg ± 14.0 mg.	PASI75 PASI90	W 24 W 48 W 108 W 156 W 204 W 264	PASI50: 67.9% PASI75: 38.1% PASI90: 14.9% PASI75: 36.6% PASI75: 40.8% PASI75: 50% PASI75: 59.4% PASI75: 60%
Menting (18), 2014 (n=178 TE ^a)	Prospl.	Etanercept initiated by 50 mg BIW SC (W0 – 12) or 50 mg once W or 25 mg BIW. Thereafter, 50 mg BIW or dose frequency reduction (every 5, 7 or 10 days).	PASI75 PASI90 Naive PASI75 PASI90 Non-naive	W 12 W 12 W 52 W 12 W 52	PASI75: 32% PASI90: 9.9% PASI90: 13.3% PASI75: 12% PASI90: 4% PASI90: 13%
Infliximab					
Kalb (41), 2005 (n=52 ^a)	Retrospl.	Infliximab 5 mg/kg IV. W 0, 2, 6 and every 8 W.	PhGA 0 PhGA 1	W 14 (licensed dose)	PhGA 0: 38.5% PhGA 1: 50%
Gisondi (40), 2008 (n=40)	Retrospl.	Infliximab 5 mg/kg W 0, 2, 6 and every 8 W.	MPD	W 24 (licensed dose)	MPD: 88.8%
Gisondi (36), 2013 (n=83)	Prospl.	Infliximab 5 mg/kg W 0, 2, 6 and every 8 W.	PASI50 PASI75 MPD	W 4 (licensed dose) W 28 (licensed dose)	PASI75: 32% PR: 64% PASI50: 96% PASI75: 69% MPD: 85%
Menting (18), 2014 (n=40 TE ^a)	Prospl.	Infliximab ± 5 mg/kg W 0, 2, 6 and every 8 W.	PASI75 PASI90 Naive	W 12 W 52	PASI75: 53% PASI90: 35.3% PASI90: 18.2%

<i>Ustekinumab</i>					
Clemmensen (37), 2011 (n=71)	Prospr.	Ustekinumab 45 or 90 mg (W 0, 4, and every 12 W). ^e	PASI75	W 12	PASI75: 38% PASI90: 12% PASI90: 14%
Laws (38), 2012 (n=129 ^a)	Retrospr.	Ustekinumab 45 or 90 mg (W 0, 4 and every 12 W). ^e	PASI50 PASI75 PASI90	W 16 (licensed dose) W 16 (licensed dose)	PASI75: 80% PASI50: 82.7% PASI75: 63% PASI90: 29.1% PASI75: 66.7% PASI75: 65.5%
Ruiz Salas (39), 2012 (n=36 ^a)	Retrospr.	Ustekinumab 45 mg (W 0, 4 and every 8–12 W). ^e	PASI50 PASI75 PASI90 MPD	W 24 (licensed dose) W 48 (licensed dose) W 12	PASI50: 89.6% PASI75: 79.3% PASI90: 65.5% PR: 80.7% PASI50: 89.6% PASI75: 79.3% PASI90: 75.9% MPD: 80.9%
		Ustekinumab 90 mg (W 0, 4 and every 8–12 W).	PASI50 PASI75 PASI90 MPD	W 24 W 36	PASI50: 89.6% PASI75: 89.6% PASI90: 75.9% MPD: 89.3% PASI50: 85.7% PASI75: 57.1% PASI90: 28.5% MPD: 75.6% PASI50: 57.1% PASI75: 28.5% PASI90: 14.2% MPD: 48.2% PASI50: 85.7% PASI75: 71.4% PASI90: 54.1% MPD: 80% PASI50: 91.7% PASI75: 75% PASI90: 58.3% MPD: 79.5% PASI50: 83.3% PASI75: 69.4% PASI90: 63.9% MPD: 71.9% PASI50: 88.8% PASI75: 71.4% PASI90: 57.1% MPD: 86.8%
Gisondi (36), 2013 (n=79)	Prospr.	Ustekinumab 45 or 90 mg (W 0, 4 and every 12 W). Dosing is weight dependent. Individuals weight (\leq 100 kg= ustekinumab 45mg; >100kg= 90 mg).	PASI50 PASI75 PASI90 MPD	W 12 W 24 W 36	PASI50: 28% MPD: 82% PASI50: 82% PASI75: 58% MPD: 60%
<i>Acitretin</i>					
Piaserico (42), 2014 (n=62)	Retrospr.	Acitretin mean 0.38 mg/kg/day.	PASI75	W 12	PASI75: 27%
<i>Fumarates</i>					
Carboni (43), 2004 (n=40)	Prospr.	Fumarates 30–360 mg/day. By remission doses were gradually reduced.	MPD	W 4 W 12 W 24	MPD: 39.6% MPD: 63% MPD: 80%
Inzinger (44), 2013 (n=272)	Retrospr.	Fumarates 30–720 mg/day.	PASI50 PASI75 PASI90 PASI100	W 12 W 24	PASI50: 76% PASI75: 47% PASI90: 9% PASI100: 1% PASI50: 82% PASI75: 63% PASI90: 27% PASI100: 5%

			W 48	PASI50: 90% PASI75: 76% PASI90: 34% PASI100: 12%
Cyclosporine				
Piaserico (42), 2014 Retrosp. Cyclosporine mean 3.5 mg/kg/day. (n=36)	PASI75	W 12		PASI75: 46%
Methotrexate				
Gisondi (40), 2008 Retrosp. Methotrexate 15 mg/week IM. (n=43)	MPD	W 24		MPD: 47.6%
Inzinger (44), 2013 Retrosp. Methotrexate 10 mg/week SC increased to 20 mg/week SC. (n=72)	PASI50 PASI75 PASI90 PASI100	W12 W 24		PASI50: 67% PASI75: 40% PASI90: 12% PASI100: 10% PASI50: 76% PASI75: 62% PASI90: 28% PASI100: 10%
Piaserico (42), 2014 (n=74) Retrosp. Methotrexate mean 11.7 mg/week.	PASI75	W 12	W 48	PASI50: 87.5% PASI75: 81% PASI90: 44% PASI100: 19% PASI75: 49%

^aAdditional conventional systemic therapies. ^bPatient switched from etanercept. ^cPatients with psoriasis with or without psoriatic arthritis. ^dPatients switching from efalizumab. ^eDosing is weight dependent. Individuals weight (≤ 100 kg= ustekinumab 45 mg; > 100 kg= 90 mg).
Retrosp.: retrospective; Prospl.: prospective; BIW: biweekly; BSA: body surface area; MBSAD: mean BSA decrease; EOW: every other week; IM: intramuscular; IV: intravenous; PASI: Psoriasis Area and Severity Index; MPD: mean PASI decrease; PhGA: Physician Global Assessment; MPhGAD: mean PhGA decrease; SC: subcutaneous; TE: treatment episode.