Supplementary material to article by S. Moreno et al. "Epidemiology of Melanocytic Naevi in Children from Lleida, Catalonia, Spain: Protective Role of Sunscreen in the Development of Acquired Moles"

Table SIV. Sunscreen use stratified by age in the sample of 4-, 8- and 14-year-old children

Sunscreen use	Age			
	4 years	8 years	14 years	Differences
Always/almost	37 (20.3%)	13 (10.4%)	4 (6.5%)	χ^2 test $p = 0.007289$, where: OR (8 vs. 4)=0.46, 95% CI=[0.21, 0.93] OR (14 vs. 8)=0.60, 95% CI=[0.14, 2.04]
Always/almost at beach/pool	146 (80.2%)	100 (80.0%)	37 (59.7%)	χ^2 test p = 0.002389, where: OR (8 vs. 4) = 0.99, 95% CI = [0.54, 1.83] OR (14 vs. 8) = 0.37, 95% CI = [0.18, 0.77]
Always/almost SPF > 15 and every 2 h	28 (15.4%)	4 (3.2%)	1 (1.6%)	72 test p=0.000100, where: OR (8 vs. 4)=0.18, 95% CI=[0.05, 0.54] OR (14 vs. 8)=0.50, 95% CI=[0.01, 5.17]
Always/almost SPF $>$ 15 and every 2 h at beach/pool	94 (51.6%)	53 (42.4%)	21 (33.9%)	χ^2 test p = 0.036166, where: OR (8 vs. 4) = 0.69, 95% CI = [0.42, 1.12] OR (14 vs. 8) = 0.70, 95% CI = [0.35, 1.37] OR (14 vs. 4) = 0.48, 95% CI = [0.25, 0.91]

SPF: sun protection factor; OR: odds ratio; 95% CI: 95% confidence interval.