

Table SI. Relative frequency of cutaneous lymphoma (CL) in Korea compared with previous studies

	This study		JSCS (8)	DACLG	SEER (4)	Previous Korea	Previous Korea
	CL %	PCL %	PCL %	(1) PCL %	PCL %	(single-centre) (3) PCL, %	(multicentre) (7) CL, %
Mature T-cell and NK-cell lymphomas	81.3	88	85.7	77	71.3	84.3	96.2
Mycosis fungoides (MF)	40	49	43.3	47	38.3	21.9	42.5
Folliculotropic MF	0.7	0.9	–	4	–	1	–
Pagetoid reticulosis	0.0	0.0	–	<1	–	–	–
Granulomatous slack skin	0.2	0.3	–	<1	–	–	–
Sézary syndrome	0.5	0.0	1.9	3	0.8	0	–
Adult T-cell leukaemia/lymphoma	0.0	0.0	16.7	–	0.1	0	–
Cutaneous CD30 ⁺ lymphoproliferative disorders	17.3	20.7	12.0	20	10.2	–	–
Lymphomatoid papulosis	7.6	9.6	3.8	12	–	9.4	5
Cutaneous anaplastic large cell lymphoma	9.7	11.1	7.8	8	–	13.5	19
Subcutaneous panniculitis-like T-cell lymphoma	3.6	4.5	2	1	0.6	10.4	11
EBV-associated NK/T-cell lymphoma	9	6	2.3	–	–	–	–
Extranodal NK/T-cell lymphoma, nasal-type	8.5	5.4	2.3	<1	0.3	16.7	15
Hydroa vacciniforme-like T-cell lymphoma	0.5	0.6	–	–	–	–	–
Cutaneous peripheral T-cell lymphoma, unspecified	10.5	7.8	5.8	5	20.8	10.4	7.5
Cutaneous CD8 ⁺ aggressive epidermotropic cytotoxic T-cell lymphoma (provisional)	0.5	0.6	0.4	<1	–	–	–
Cutaneous γ/δ^+ T-cell lymphoma (provisional)	1.7	1.8	0.3	<1	–	2.1	–
Cutaneous CD4 ⁺ small/medium-sized pleomorphic T-cell lymphoma (provisional)	2.1	2.7	1.4	2	–	5.2	–
Angioimmunoblastic T-cell lymphoma	1.9	0.0	–	–	0.2	0	1
Mature B-cell lymphomas	17.8	11.7	12.9	23	28.5	13.5	3.8
Cutaneous marginal zone B-cell lymphoma	3.8	4.5	4.2	7	7.1	9.4	1
Cutaneous follicle centre B-cell lymphoma	1	0.9	2.1	11	8.5	0	3
Cutaneous diffuse large B-cell lymphoma	11.9	6.3	–	–	11.4	–	–
Primary cutaneous diffuse large B-cell lymphoma, leg-type	4.5	5.7	5.5	4	2.6	1	–
Primary cutaneous diffuse large B-cell lymphoma, other	0.5	0.6	–	<1	8.8	2.1	–
Intravascular large B-cell lymphoma	0.5	0.0	1.2	<1	–	1	–
Burkitt lymphoma	–	0.2	0.0	–	0.1	0	0
Precursor haematological neoplasm	0.9	0.3	1.4	–	0.3	2	–
CD4 ⁺ /CD56 ⁺ haematodermic neoplasm (formerly blastic NK cell lymphoma)	0.9	0.3	1.4	–	0.2	–	–
Hodgkin's lymphoma	0.5	0.0	–	–	–	–	–
Total	422	333	1,733	1,905	3,884	164	80

PCL: primary cutaneous lymphoma; JSCS: Japanese Skin Cancer Society; DACLG: Dutch and Austrian Cutaneous Lymphoma Group; SEER: surveillance, epidemiology, and end results; NK: natural killer; MF: mycosis fungoides; EBV: Epstein-Barr virus.