

Table SII. Pattern of long-term sequelae in 102 patients with Stevens-Johnson syndrome (SJS)/toxic epidermal necrolysis (TEN) overlap, and TEN

Sequelae	All cases n=102 n (%)	SJS n=81 n (%)	SJS/TEN overlap + TEN n=21 n (%)	p-value
All sequelae	68 (66.7)	51 (63.0)	17 (81.0)	0.119
<i>Cutaneous</i>				
Chronic eczema	45 (44.1)	35 (43.2)	10 (47.6)	0.887
Nail change	32 (31.4)	24 (29.6)	8 (38.1)	0.456
Post-inflammatory hypo- or hyper-pigmentation	12 (11.8)	6 (7.4)	6 (28.6)	0.007
Vitiligo	14 (13.7)	11 (13.6)	3 (14.3)	0.933
Autoimmune bullous disease	2 (2.0)	1 (1.2)	1 (4.8)	0.299
<i>Ocular</i>				
Dry eye syndrome	22 (21.6)	15 (18.5)	13 (61.9)	0.065
Chronic conjunctivitis	6 (5.9)	1 (1.2)	9 (42.9)	0.248
Trichiasis	5 (4.9)	4 (4.9)	1 (4.8)	0.973
Corneal erosion	4 (3.9)	1 (1.2)	3 (14.3)	0.006
Symblepharon/Synechia	2 (2.0)	1 (1.2)	1 (4.8)	0.299
Meibomian gland dysfunction	1 (1.0)	1 (1.2)	0 (0.0)	0.609
Nasolacrimal duct obstruction	1 (1.0)	1 (1.2)	0 (0.0)	0.609
<i>Connective tissue diseases</i>				
Systemic lupus erythematosus	2 (2.0)	1 (1.2)	0 (0)	0.609
Hashimoto thyroiditis	1 (1.0)	1 (1.2)	0 (0)	0.609
Primary Sjögren's syndrome	1 (1.0)	2 (2.5)	0 (0)	0.467
<i>Visceral organ involvement</i>				
Pulmonary	2 (2.0)	1 (1.2)	0 (0)	0.609
Renal	1 (1.0)	2 (2.5)	0 (0)	0.467

Significant values are shown in bold.