

Table SII. Brief Pain Inventory (BPI) questions among patients with hidradenitis suppurativa (HS) according to Hurley stage of HS severity

11-point numerical rating scale	Hurley stage I	Hurley stage II	Hurley stage III	All patients
Age, years, median (range)	38 (24–48)	44 (23–63)	46 (33–66)	44 (35–51)
Patients, <i>n</i>	7	29	9	45
Sex (female:male), <i>n</i>	6:1	23:6	4:5	33:12
Sartorius	9 (7–11)	20 (13–27)	44 (41–91)*, **	20 (11–38)
Pain severity questions				
Pain at its worst in the last 24 h	0 (0–1.75)	4 (1.5–7)	7 (4.5–8.5)*	4 (0–7)
Pain at its least in the last 24 h	0 (0–0)	2 (0.5–5)	4 (1.5–5.25)*	2 (0–5)
Pain at its average in the last 24 h	1 (0.25–2)	4 (1.5–6.5)	6 (3–8)*	4 (1–6.25)
Pain at the time of investigation	0 (0–0)	2.5 (0–5.5)	6 (3.75–8.25)*	2 (0–6)
Pain relief of subscribed medicine (%)	40 (0–90)	20 (20–70)	30 (7.5–65)	10 (0–30)
Pain interference, median (range)				
General activity	0 (0–0)	2 (0–6)	7 (2.25–8.25)*	2 (0–6)
Mood	2 (0–2.75)	4 (1.25–6)	4 (1.5–6.5)	3 (1–6)
Walking ability	0 (0–0.75)	2 (0–5.75)	5 (2.25–9)	2 (0–6)
Ability to work	0 (0–0)	2 (0–5.75)	5 (0.75–7)*	2 (0–5.5)
Relationships with other people	0 (0–0)	4 (0.25–6)	5 (4–6.25)*	3 (0–5.5)
Sleep	0 (0–0)	3 (0–5)	3 (1–8.5)*	2 (0–5)
Enjoyment of life	0 (0–0.75)	4 (1.25–6.75)	4 (3.5–9)*	4 (0–7)
Anxiety score (HAD-A), median (range)	7 (2.75–12.25)	9.5 (6.5–12.5)	7 (3.75–15.25)	9 (5–14)
Depression score (HAD-D), median (range)	2 (1–8.25)	9 (2.5–11.5)	11 (3.5–13.5)	9 (2–12)

Data are presented as median (interquartile range), except for age.

**p*-values <0.05 between Hurley stages I and III; and **between Hurley stages II and III.