

Table SI. Characteristics of patients with Bowen's disease (BD) according to their immunosuppression status

Variable	Non-immunocompromised (n=79)	Immunocompromised (n=25)	p-value
Median age at first PDT, years (IQR)	77 (66–82)	69 (61–79)	0.11
Sex, n (%)			0.35
Women	50 (63)	13 (52)	
Men	29 (37)	12 (48)	
History of SCC, n (%)			0.004
No	55 (73)	10 (40)	
Yes	20 (27)	15 (60)	
NA	4	0	
More than one PDT field, n (%)			0.007
No	65 (82)	13 (52)	
Yes	14 (18)	12 (48)	
At least one BD on the lower limbs, n (%)			0.62
No	54 (68)	19 (76)	
Yes	25 (32)	6 (24)	
At least one BD on the upper limbs, n (%)			0.071
No	61 (77)	14 (56)	
Yes	18 (23)	11 (44)	
At least one BD on the head or neck, n (%)			0.65
No	37 (47)	10 (40)	
Yes	42 (53)	15 (60)	
At least one BD on the trunk, n (%)			0.13
No	68 (86)	18 (72)	
Yes	11 (14)	7 (28)	
Minimal number of treated BD, median (IQR)	1 (1–1)	1 (1–1)	0.67
Not available, n	2	0	
Minimal number of session/PDT field, median (IQR)	2 (2–2)	2 (2–2)	0.003

PDT: photodynamic therapy; SCC: squamous cell carcinoma.