

Table SI. Recent studies of 1-year mortality for patients with bullous pemphigoid (search terms bullous pemphigoid and mortality in PubMed, written in English, since 1990)

First author and year of publication (ref)	Country/region	Study period (duration, years)	Cases, <i>n</i>	1-year mortality, %	Standardized mortality ratio
Venning 1992 (11)	UK	1975–1988 (13)	82	19	–
Bernard 1997 (12)	France	1987–1994 (7)	94	37	–
Roujeau 1998 (13)	France	1985–1992 (7)	217	41	–
Rzany 2002 (14)	Germany	1987–1997 (10)	369	29	–
Colbert 2004 (15)	USA	1997–2002 (5)	37	11	–
Gudi 2005 (7)	Scotland	1991–2001 (10)	83	25	5.8
Joly 2005 (16)	France	1996–1998 (2)	341	26	–
Langan 2008 (6)	UK	1996–2006 (10)	869 ^a	19	2.3
Parker 2008 (17)	USA	1998–2003 (5)	223	23	0.4–0.7
Cortés 2011 (18)	Switzerland	2001–2004 (3)	115	21	3.0
Cortés 2012 (19)	Switzerland	1990–2003 (13)	60	27	–
Joly 2012 (12)	France	2000–2005 (5)	312	38	6.6
Cai 2013 (20)	Singapore	2004–2009 (5)	359	27	2.7 (3-year ^c)
Li 2013 (21)	China	1991–2011 (20)	140	13	3.1–6.1
Zhang 2013 (22)	China	2005–2010 (5)	94	23	–
Brick 2014 (1)	USA	1960–2009 (49)	87	19	1.9
Försti 2015 ^b	Finland	1985–2012 (27)	198	17	7.6
Gual 2014 (23)	Spain	1990–2010 (20)	101	13	2.3
Lee 2014 (39)	Korea	1993–2013 (20)	168	19	2.4–9.6
Serwin 2014 (40)	Poland	1999–2012 (13)	122	25	–

^aDiagnoses based on general practice database without immunological confirmation. ^bCurrent study. ^cNot for 1 year as conventionally.