Supplementary material to article by A. Batycka-Baran et al. "Decreased Number of Circulating Endothelial Progenitor Cells (CD133<sup>+</sup>/KDR<sup>+</sup>) in Patients with Psoriatic Arthritis"

 $\label{thm:characteristics} Table\,SI.\,\textit{Clinical and demographic characteristics of patients with psoriatic arthritis\,(PsA)\,(n=24)\,and\,psoriasis\,vulgaris\,(n=63)\,(22)\,$ 

Characteristics	Patients with PsA	Patients with plaque psoriasis $(n=63)$	р
Age, years, mean ± SD Range	48.29 ± 9.05 26–66	42.16 ± 15.23 18–73	0.063
Male/female, %	62.5/37.5	51.6/44.5	0.74
Obesity <sup>a</sup> , n (%)	6 (25.5)	11 (17.46)	0.32
Hypertension <sup>b</sup> , n (%)	7 (29.1)	10 (15.87)	0.16
Smoking habit <sup>c</sup> , n (%)	9 (37.5)	28 (44.4)	0.16

<sup>&</sup>lt;sup>a</sup>Body mass index ≥30. <sup>b</sup>Systolic blood pressure ≥140 mmHg or diastolic blood pressure ≥90 mmHg or use of antihypertensive medication. <sup>c</sup>Smoking at least 1 cigarette per day during the last year.