

Table SII. Comparison of relative frequency of cutaneous lymphoma between subgroups per 10-year period between 1994 and 2013

	SCL, n (%)		PCL, n (%)	
	1994 to 2003 n = 33	2004 to 2013 n = 73	1994 to 2003 n = 102	2004 to 2013 n = 187
<i>Mature T- and NK-cell lymphoma</i>	25 (75.8)	34 (46.6)	92 (90.2)	152 (81.3)
Mycosis fungoides	0 (0)	0 (0)	27 (26.5)	58 (31.0)
Lymphomatoid papulosis	0 (0)	0 (0)	21 (20.6)	25 (13.4)
Cutaneous anaplastic large cell lymphoma	5 (15.2)	6 (8.2)	9 (8.8)	25 (13.4)
Subcutaneous panniculitis-like T-cell lymphoma	0 (0)	0 (0)	7 (6.9)	14 (7.5)
Extranodal NK/T-cell lymphoma	9 (27.3)	11 (15.1)	14 (13.7)	14 (7.5)
Peripheral T-cell lymphoma, unspecified	11 (33.3)	11 (15.1)	14 (13.7)	13 (7.0)
Cutaneous γ/δ^+ T-cell lymphoma			0 (0)	0 (0)
Cutaneous CD4 ⁺ small/medium pleomorphic T-cell lymphoma			0 (0)	3 (1.6)
Angioimmunoblastic lymphoma	0 (0)	5 (6.8)	0 (0)	0 (0)
Adult T-cell lymphoma/leukaemia	0 (0)	1 (1.4)	0 (0)	0 (0)
<i>Mature B-cell lymphoma</i>	5 (15.2)	32 (43.8)	8 (7.8)	32 (17.1)
Cutaneous marginal zone B-cell lymphoma (MALT lymphoma)	0 (0)	1 (1.4)	5 (4.9)	19 (10.2)
Cutaneous follicle centre lymphoma	0 (0)	1 (1.4)	0 (0)	1 (0.53)
Cutaneous diffuse large B-cell lymphoma, leg type	0 (0)	0 (0)	3 (2.9)	11 (5.9)
Cutaneous diffuse large B-cell lymphoma, others	4 (12.1)	26 (35.6)	0 (0)	0 (0)
Intravascular large B-cell lymphoma	0 (0)	1 (1.4)	0 (0)	1 (0.53)
Burkitt's lymphoma	1 (3.0)	2 (2.7)	0 (0)	0 (0)
Mantle cell lymphoma	0 (0)	1 (1.4)	0 (0)	0 (0)
<i>Immature haematopoietic malignancies</i>	2 (6.1)	7 (9.6)	2 (2.0)	3 (1.6)
Lymphoblastic lymphoma	2 (6.1)	5 (6.8)	1 (0.98)	2 (1.0)
Blastic plasmacytoid dendritic cell neoplasm	0 (0)	2 (2.7)	1 (0.98)	1 (0.53)

SCL: secondary cutaneous lymphoma; PCL: primary cutaneous lymphoma.