

Table SIII. Clinical features of the skin lesions in primary cutaneous lymphoma patients in this study

	Site of cutaneous involvement, n (%)					Morphology of skin lesions, n (%)					Multiple skin lesions n=202 n (%)
	Head and neck n=82	Trunk n=141	Arms n=113	Legs n=132	Whole body n=44	Nodule n=119	Papule n=75	Plaque n=45	Maculo-patch n=83	Cellulitis-like swelling or ulceration n=25	
Mature T- and natural killer-cell lymphoma (n=244)	58 (23.8)	134 (54.9)	108 (44.3)	124 (50.8)	43 (17.6)	82 (33.6)	69 (28.3)	42 (17.2)	81 (33.2)	23 (9.4)	190 (77.9)
Mycosis fungoides (n=85)	7 (8.2)	59 (69.4)	37 (43.5)	41 (48.2)	23 (27.1)	8 (9.4)	9 (10.6)	28 (32.9)	61 (71.8)	0 (0)	79 (92.9)
Lymphomatoid papulosis (n=46)	8 (17.4)	27 (58.7)	24 (52.2)	32 (69.6)	15 (32.6)	12 (26.1)	42 (91.3)	6 (13.0)	8 (17.4)	0 (0)	42 (91.3)
Anaplastic large cell lymphoma (n=34)	11 (32.4)	10 (29.4)	14 (41.2)	14 (41.2)	2 (5.9)	22 (64.7)	10 (29.4)	1 (2.9)	0 (0)	1 (2.9)	16 (47.1)
Subcutaneous panniculitis-like T-cell lymphoma (n=21)	6 (28.6)	12 (57.1)	10 (47.6)	10 (47.6)	0 (0)	15 (71.4)	0 (0)	4 (19.0)	4 (19.0)	3 (14.3)	17 (80.9)
Extranodal natural killer/T-cell lymphoma (n=28)	9 (32.1)	13 (46.4)	12 (42.3)	18 (64.3)	0 (0)	12 (42.9)	0 (0)	0 (0)	0 (0)	18 (64.3)	19 (67.9)
Peripheral T-cell lymphoma, unspecified (n=27)	15 (55.6)	13 (48.1)	10 (37.0)	9 (33.3)	3 (11.1)	12 (44.4)	7 (25.9)	3 (11.1)	7 (25.9)	1 (3.7)	15 (55.6)
Cutaneous γ/δ + T-cell lymphoma (n=0)	a	a	a	a	a	a	a	a	a	a	a
CD4+ small/medium pleomorphic T-cell lymphoma (n=3)	2 (66.7)	0 (0)	1 (33.3)	0 (0)	0 (0)	1 (33.3)	1 (33.3)	0 (0)	1 (33.3)	0 (0)	2 (66.7)
Mature B-cell lymphoma (n=40)	21 (52.5)	7 (17.5)	4 (10.0)	8 (20.0)	0 (0)	32 (80.0)	6 (15.0)	3 (7.5)	2 (5.0)	2 (5.0)	11 (27.5)
Marginal zone B-cell lymphoma (n=24)	18 (75)	3 (12.5)	3 (12.5)	1 (4.2)	0 (0)	20 (83.3)	2 (8.3)	1 (4.2)	1 (4.2)	0 (0)	2 (8.3)
Follicle centre lymphoma (n=1)	1 (100)	0 (0)	0 (0)	0 (0)	0 (0)	1 (100)	0 (0)	0 (0)	0 (0)	0 (0)	1 (100)
Diffuse large B-cell lymphoma, leg type (n=14)	2 (14.3)	4 (28.6)	1 (7.1)	6 (42.9)	0 (0)	11 (78.6)	4 (28.6)	2 (14.3)	0 (0)	2 (14.3)	7 (50)
Diffuse large B-cell lymphoma, others (n=0)	a	a	a	a	a	a	a	a	a	a	a
Intravascular large B-cell lymphoma (n=1)	0 (0)	0 (0)	0 (0)	1 (100)	0 (0)	0 (0)	0 (0)	0 (0)	1 (100)	0 (0)	1 (100)
Immature haematopoietic malignancies (n=5)	3 (60.0)	0 (0)	1 (20.0)	0 (0)	1 (20.0)	5 (100)	0 (0)	0 (0)	0 (0)	0 (0)	1 (20.0)
B-cell lymphoblastic lymphoma (n=3)	2 (66.7)	0 (0)	1 (33.3)	0 (0)	0 (0)	3 (100)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Blastic plasmacytoid dendritic cell neoplasm (n=2)	1 (50.0)	0 (0)	0 (0)	0 (0)	1 (50.0)	2 (100)	0 (0)	0 (0)	0 (0)	0 (0)	1 (50.0)

a)No data found.