

Table SIV. Extent of skin lesions and extracutaneous dissemination in primary cutaneous lymphoma

	T classification at initial work-up				Visceral		
	T1, N n (%)	T2 n (%)	T3 n (%)	T4 n (%)	LN involve- ment n (%)	involve- ment n (%)	Site of viscera
Mature T- and natural killer-cell lymphoma							
Mycosis fungoides (n=85)	34 (40.0)	37 (43.5)	8 (9.4)	6 (7.1)	16 (18.8)	1 (1.2)	Pleura (n=1)
Lymphomatoid papulosis (n=45)	4 (8.9)	9 (20)	33 (73.3)	^a	0 (0)	0 (0)	
Anaplastic large cell lymphoma (n=34)	18 (52.9)	4 (11.8)	12 (35.3)	^a	1 (2.9)	0 (0)	
SPTL (n=21)	3 (14.3)	1 (4.8)	17 (80.9)	^a	0 (0)	2 (9.5)	CNS (n=1), Pleura (n=1)
Extranodal NKTL (n=28)	6 (21.4)	7 (25)	15 (53.6)	^a	15 (53.6)	12 (42.9)	Gastrointestinal tract (n=8), lung (n=7), liver (n=4), soft tissue (n=3), kidney (n=3), bone (n=3), pancreas (n=3)
PTCL, unspecified (n=27)	12 (44.4)	4 (14.8)	11 (40.7)	^a	4 (14.8)	5 (18.5)	
GDTL (n=0)							Oesophagus (n=1), stomach (n=1), bone
SMTL (n=3)	1 (33.3)	1 (33.3)	1 (33.3)		0 (0)	0 (0)	(n=1), liver (n=1), lung (n=1)
Mature B-cell lymphoma							
MZBCL (n=24)	22 (91.7)	0 (0)	2 (8.3)	^a	0 (0)	0 (0)	
Follicle centre lymphoma (n=1)	0 (0)	1 (100)	0 (0)	^a	0 (0)	0 (0)	
DLBCL, leg type (n=14)	9 (64.3)	4 (28.6)	1 (7.1)	^a	8 (57.1)	3 (21.4)	Adrenal gland (n=2), brain (n=1), thyroid (n=1), spleen (n=1), pericardium (n=1)
DLBCL, others (n=0)							
ILBCL (n=1)	0 (0)	0 (0)	1 (100)	^a	0 (0)	0 (0)	
Immature haematopoietic malignancies							
B-cell lymphoblastic lymphoma (n=3)	3 (100)	0 (0)	0 (0)	^a	0 (0)	0 (0)	
BPDCN (n=2)	1 (50.0)	0 (0)	1 (50.0)	^a	1 (50.0)	0 (0)	

^aNo data found.

BPDCN: blastic plasmacytoid dendritic cell neoplasm; DLBCL: diffuse large B-cell lymphoma; GDTL: cutaneous γ/δ^+ T-cell lymphoma; ILBCL: intravascular large B-cell lymphoma; LN: lymph node; MZBCL: marginal zone B-cell lymphoma; NKTL: natural killer/T-cell lymphoma; PTCL: peripheral T-cell lymphoma; SMTL: CD4⁺ small/medium pleomorphic T-cell lymphoma; SPTL: subcutaneous panniculitis-like T-cell lymphoma.