

Table SI. Mixed effects multiple regression with a random intercept for each patient of baseline Psoriasis Area Severity Index (PASI) vs predictors and the difference in $\log(1+\text{PASI})$ between 3 months and baseline, respectively

	Baseline PASI			$\log(1+\text{PASI}_{3\text{m}})-\log(1+\text{PASI}_{\text{BL}})$			$\log(1+\text{PASI}_{6\text{m}})-\log(1+\text{PASI}_{\text{BL}})$			$\log(1+\text{PASI}_{24\text{m}})-\log(1+\text{PASI}_{\text{BL}})$		
	Coefficient estimate	SE	<i>p</i> -value	Coefficient estimate	SE	<i>p</i> -value	Coefficient estimate	SE	<i>p</i> -value	Coefficient estimate	SE	<i>p</i> -value
Intercept	1.92	0.39	< 0.0001	-0.06	0.44	0.14	-0.85	0.73	0.25	-1.03	0.66	0.12
Naïve vs not naïve (ref.)	-0.39	0.08	< 0.0001	0.39	0.13	0.005	0.66	0.19	0.004	0.50	0.17	0.023
Sex, male vs. female (ref.)	0.54	0.18	0.003	-0.11	0.20	0.56	0.06	0.33	0.86	-0.07	0.27	0.79
Age, years	0.0004	0.005	0.93	0.006	0.005	0.27	-0.004	0.009	0.61	-0.003	0.007	0.70
Body mass index, kg/m ²	0.012	0.012	0.32	-0.021	0.013	0.12	-0.007	0.019	0.72	-0.007	0.020	0.75

SE: standard error. Significant values ($p < 0.05$) are shown in **bold**.