

Table SIV. Characteristics of the 284 ulcers treated with punch-graft

Ulcers of the foot, <i>n</i> (%)	31 (10.9)
Ulcers of the leg, <i>n</i> (%)	253 (89.1)
Venous ulcers, <i>n</i> (%)	124 (43.7)
Arterial ulcers, <i>n</i> (%)	32 (11.3)
Diabetic ulcers, <i>n</i> (%)	4 (1.4)
Pressure ulcers, <i>n</i> (%)	0
Mixed ulcers, <i>n</i> (%)	68 (23.9)
Other ulcers, <i>n</i> (%)	56 (19.7)
Vasculitis ulcer	27
Trauma-induced ulcer	14
Infection-induced ulcer	4
Pyoderma gangrenosum	3
Oedema-induced ulcer (without venous insufficiency)	2
Pharmaceutical side-effect-induced ulcer	2
Radiodermatitis-induced ulcer	2
Fistula ulcer (without known cause)	1
Necrobiosis lipoidica	1
Duration of ulcer presence at baseline, months, mean ± SD	24.9 ± 33.8
Venous ulcers	30.9 ± 37.9
Arterial ulcers	20.6 ± 23.2
Diabetic ulcers	21.0 ± 17.2
Mixed ulcers	22.2 ± 23.5
Other ulcers	18.7 ± 39.6
Duration of ulcer ≥ 6 months, <i>n</i> (%)	193 (68.0)

SD: standard deviation.