

Table SI. Tumour, nodes, metastases, blood (TNMB) classification (International Society for Cutaneous Lymphomas/European Organization of Research and Treatment of Cancer (ISCL/EORTC) revision (21))

TNMB classification		
Skin	T ₁	Patches ¹ , papules and/or plaques ² <10% of body surface (T _{1a} : Patch – T _{1b} : Plaque +/- patch)
	T ₂	Patches, papules and/or plaques >10% of body surface (T _{2a} : Patch – T _{2b} : Plaque +/- patch)
	T ₃	One or more tumours (>1 cm diameter)
	T ₄	Confluent erythema >80% of body surface
Node	N ₀	No clinically abnormal peripheral lymph nodes
	N ₁	Clinically abnormal peripheral lymph nodes ³ ; histopathology Dutch grade 1 or NCI LN0-2. (N _{1a} : clone-, N _{1b} : clone+)
	N ₂	Clinically abnormal peripheral lymph nodes; histopathology Dutch grade 2 or NCI LN3 (N _{2a} : clone-, N _{2b} : clone +)
Visceral	M ₀	No visceral organs involved
	M ₁	Viscera involved (with pathological confirmation – imaging for spleen/liver)
Blood	B ₀	<5% of peripheral blood lymphocytes are Sézary cells ⁴ (B _{0a} : clone- B _{0b} : clone +)
	B ₁	>5% of peripheral blood lymphocytes are Sézary cells (B _{1a} : clone- B _{1b} : clone +)
	B ₂	High tumour burden: ≥ 1,000/μl Sézary cells with clone +

¹patches- not significantly elevated nor indurated skin lesions of any size; ²plaques- elevated or indurated skin lesions of any size, ³abnormal peripheral lymph nodes - firm, irregular, clustered, fixed or >=1.5 cm in diameter lymph nodes. ⁴Sézary cells - atypical lymphocytes with hyperconvoluted, cerebriform nuclei