Table SII. Follow-up time, number of events, incidence rates, and hazard ratios (HR) of myocardial infarction, ischaemic stroke, cardiovascular death, and major adverse cardiovascular events, respectively

	Chronic urticaria		Inducible urticaria	
	Controls	Cases	Controls	Cases
Myocardial infarction				
Follow-up time, years	385,254	13,102	184,850	6,286
Number of events	568	25	223	9
Incidence rate/1,000 person-years (95% confidence interval)	1.47 (1.36-1.60)	1.91 (1.29-2.82)	1.21 (1.06-1.38)	1.43 (0.75-2.75)
Unadjusted HR (95% confidence interval)	1.29 (0.87-1.93, p=0.21)		1.19(0.61-2.31, p=0.62)	
Adjusted HR (95% confidence interval, p-value)	1.18 (0.79–1.76, p=0.42)		1.07 (0.55-2.08, p=0.85)	
Ischaemic stroke				
Follow-up time, years	384,578	13,059	184,784	6,282
Number of events	721	27	253	7
Incidence rate/1,000 person-years (95% confidence interval)	1.87 (1.74-2.02)	2.07 (1.42-3.01)	1.37 (1.21-1.55)	1.11 (0.53-2.34)
Unadjusted HR (95% confidence interval)	1.10 (0.75-1.62, <i>p</i> =0.62)		0.81 (0.38–1.72, <i>p</i> =0.59)	
Adjusted HR (95% confidence interval, p-value)	1.03 (0.70-1.52, p=0.88)		0.80 (0.38-1.70, p=0.57)	
Cardiovascular death				
Follow-up time, years	386,965	13,152	185,666	6,308
Number of events	526	13	153	4
Incidence rate/1,000 person-years (95% confidence interval)	1.36 (1.25-1.48)	1.00 (0.57-1.70)	0.82 (0.70-0.97)	0.63 (0.24-1.69)
Unadjusted HR (95% confidence interval)	0.73 (0.42-1.26, p=0.26)		0.77 (0.28-2.07, p=0.60)	
Adjusted HR (95% confidence interval, p-value)	0.67 (0.39-1.17, p=0.16)		0.63 (0.23-1.72, p=0.37)	
Major adverse cardiovascular events				
Follow-up time, years	382,956	13,013	183,999	6,260
Number of events	1.514	57	543	16
Incidence rate/1,000 person-years (95% confidence interval)	3.95 (3.76-4.16)	4.38 (3.38-5.68)	2.95 (2.71-3.21)	2.56 (1.57-4.17)
Unadjusted HR (95% confidence interval)	1.11(0.86-1.44, p=0.45)		0.87 (0.53-1.42, p=0.57)	
Adjusted HR (95% confidence interval, p-value)	1.09(0.83-1.42, p=0.54)		0.76(0.46-1.25, p=0.27)	

Acta Derm Venereol 2017