

Table SI. Subject characteristics of patients with bullous pemphigoid (BP) (n = 25)

Patient	Age, years/ Sex	ELISA (U/ml)		Staining		Eosinophils	
		BP180	BP230	DIF	IIF	Peripheral blood (%)	×10 ⁶ /ml BF
BP - 1	93/F	221	neg.	C3c, IgG lin. at DEJ	IgG lin.	18.0	1.4
BP - 2	95/F	345	116	C3c, IgG lin. at DEJ	IgG lin.	0.5	n.a.
BP - 3	79/M	267	148	C3c, IgG lin. at DEJ	IgG lin.	11.3	1.1
BP - 5	82/F	121	70	IgG lin. at DEJ	n.a.	n.a.	5.4
BP - 6	93/M	801	80	IgG lin. at DEJ	n.a.	0.9	n.a.
BP - 7	83/F	13	132	C3c, IgG lin. at DEJ	IgG lin.	5.0	0.6
BP - 8	82/F	115	23	C3c, IgG lin. at DEJ	IgG lin.	4.6	0
BP - 9	78/F	3	9	C3c, IgG lin. at DEJ	IgG + IgA lin.	3.1	0.5
BP - 10	64/M	118	2	C3c, IgG lin. at DEJ	IgG lin.	9.7	0.5
BP - 11	80/F	132	92	C3c, IgG lin. at DEJ	IgG lin.	0.9	1.8
BP - 13	88/F	neg.	neg.	C3c, IgG lin. at DEJ	IgG lin.	0.5	n.a.
BP - 14	71/M	65	27	IgG lin. at DEJ	IgG lin.	n.a.	8.7
BP - 15	81/F	37	84	C3c, IgG lin. at DEJ	IgG lin.	7.3	n.a.
BP - 16	92/F	421	139	n.a.	neg.	0	1.4
BP - 17	93/F	201	466	C3c, IgG lin. at DEJ	IgG+IgA lin.	13.2	9.6
BP - 18	75/F	114	42	C3c, IgG lin. at DEJ	IgG lin.	30.1	n.a.
BP - 19	61/F	174	79	C3c, IgG lin. at DEJ	IgG lin.	11.0	n.a.
BP - 20	95/F	5	11	IgM lin. at DEJ	IgG lin.	14.5	n.a.
BP - 21	73/F	129	4	n.a.	n.a.	16.1	n.a.
BP - 22	79/F	110	92	C3c, IgG lin. at DEJ	IgG lin.	9.5	n.a.
BP - 23	88/F	36	0	C3c, IgG lin. at DEJ	IgG lin.	10.4	2.6
BP - 24	86/M	196	96	C3c, IgG lin. at DEJ	IgG lin.	19.0	1.5
BP - 25	63/M	79	4	C3c, IgG lin. at DEJ	IgG lin.	4.8	n.a.

n.a.: not assessed; neg.: negative; DIF: direct immunofluorescence; IIF: indirect immunofluorescence (with monkey-oesophagus); BF: blister fluid; lin.: linear; DEJ: dermo-epidermal junction zone; ELISA: enzyme-linked immunoassay.