

**Appendix S1.****COMPLETE REFERENCE LIST**

1. Baroni A, Staibano S, Russo T, Piccolo V, Satriano RA, Vozza A, et al. Verrucous epidermal naevus and naevus spilus associated with lower limb asymmetry and right bundle-branch block: a case of phacomatosis pigmentokeratotica? *Clin Exp Dermatol* 2012; 37: 74–75.
2. Tavoloni Braga JC, Gomes E, Macedo MP, Pinto C, Duprat J, Begnami MD, et al. Early detection of melanoma arising within nevus spilus. *J Am Acad Dermatol* 2014; 70: e31–e32.
3. Sigg C, Pelloni F, Schnyder UW. Frequency of congenital nevi, nevi spili and cafe-au-lait spots and their relation to nevus count and skin complexion in 939 children. *Dermatologica* 1990; 180: 118–123.
4. Corradin MT, Zattra E, Fiorentino R, Alaibac M, Belloni-Fortina A. Nevus spilus and melanoma: case report and review of the literature. *J Cutan Med Surg* 2010; 14: 85–89.
5. Yoneyama K, Kamada N, Mizoguchi M, Utani A, Kobayashi T, Shinkai H. Malignant melanoma and acquired dermal melanocytosis on congenital nevus spilus. *J Dermatol* 2005; 32: 454–458.
6. Sarin KY, McNiff JM, Kwok S, Kim J, Khavari PA. Activating HRAS mutation in nevus spilus. *J Invest Dermatol* 2014; 134: 1766–1768.
7. Krengel S, Widmer DS, Kerl K, Levesque MP, Schiestl C, Weibel L. Naevus spilus-type congenital melanocytic naevus associated with a novel NRAS codon 61 mutation. *Br J Dermatol* 2016; 174: 642–644.
8. Perkins NG. Melanoma arising in a cafe au lait spot of neurofibromatosis. *Am J Surg* 1957; 93: 1018–1020.
9. Corradin MT, Giulioni E, Fiorentino R, Santeufemia DA, Re GL, Vettorello A. In situ malignant melanoma on nevus spilus in an elderly patient. *Acta Dermatovenerol Alp Pannonica Adriat* 2014; 23: 17–19.
10. Vion B, Belaich S, Grossin M, Preaux J. Les aspects évolutifs du naevus sur naevus: revue de la littérature à propos de 7 observations. *Ann Dermatol Venereol* 1985; 112: 813–819.
11. Stern JB, Haupt HM, Aaronson CM. Malignant melanoma in a speckled zosteriform lentiginous nevus. *Int J Dermatol* 1990; 29: 583–584.
12. Bologna JL. Fatal melanoma arising in a zosteriform speckled lentiginous nevus. *Arch Dermatol* 1991; 127: 1240–1241.
13. Borrego L, Hernandez SJ, Baez O, Hernandez HB. Naevus spilus as a precursor of cutaneous melanoma: report of a case and literature review. *Clin Exp Dermatol* 1994; 19: 515–517.
14. Breitkopf C, Ernst K, Hundeiker M. Neubildungen auf Naevus spilus. *Hautarzt* 1996; 47: 759–762.
15. Brufau C, Moran M, Armijo M. Naevus sur naevus: à propos de 7 observations, trois associées à d'autres dysplasies, et une à un mélanome. *Ann Dermatol Venereol* 1986; 113: 409–418.
16. Rhodes AR, Mihm MC, Jr. Origin of cutaneous melanoma in a congenital dysplastic nevus spilus. *Arch Dermatol* 1990; 126: 500–505.
17. Wagner RF, Jr., Cottell WI. In situ malignant melanoma arising in a speckled lentiginous nevus. *J Am Acad Dermatol* 1989; 20: 125–126.
18. Rutten A, Goos M. Nevus spilus with malignant melanoma in a patient with neurofibromatosis. *Arch Dermatol* 1990; 126: 539–540.
19. Guillot B, Bessis D, Barneon G, Monpoint S, Guilhou JJ. Malignant melanoma occurring on a 'naevus on naevus'. *Br J Dermatol* 1991; 124: 610–611.
20. Kurban RS, Preffer FI, Sober AJ, Mihm MC, Jr., Barnhill RL. Occurrence of melanoma in "dysplastic" nevus spilus: report of case and analysis by flow cytometry. *J Cutan Pathol* 1992; 19: 423–428.
21. Krahn G, Thoma E, Peter RU. Zwei superfiziell spreitende maligne Melanome auf Naevus spilus. *Hautarzt* 1992; 43: 32–34.
22. Vazquez-Doval J, Sola MA, Contreras-Mejuto F, Redondo P, Soto J, Quintanilla E. Malignant melanoma developing in a speckled lentiginous nevus. *Int J Dermatol* 1995; 34: 637–638.
23. Grinspan D, Casala A, Abulafia J, Mascotto J, Allevato M. Melanoma on dysplastic nevus spilus. *Int J Dermatol* 1997; 36: 499–502.
24. Cox NH, Malcolm A, Long ED. Superficial spreading melanoma and blue naevus within naevus spilus – ultrastructural assessment of giant pigment granules. *Dermatology* 1997; 194: 213–216.
25. Weinberg JM, Schutzer PJ, Harris RM, Tangoren IA, Sood S, Rudolph RI. Melanoma arising in nevus spilus. *Cutis* 1998; 61: 287–289.
26. Ankit C, Khirwadkar N, Azurdia RM. Malignant melanoma arising in a nevus spilus. *Dermatol Online J* 2011; 17: 10.
27. Abecassis S, Spatz A, Cazeneuve C, Martin-Villepou A, Clerici T, Lacour JP, et al. Mélanome sur naevus spilus : 5 observations. *Ann Dermatol Venereol* 2006; 133: 323–328.
28. Ly L, Christie M, Swain S, Winship I, Kelly JW. Melanoma(s) arising in large segmental speckled lentiginous nevi: a case series. *J Am Acad Dermatol* 2011; 64: 1190–1193.
29. Piana S, Gelli MC, Grenzi L, Ricci C, Gardini S, Piana S. Multifocal melanoma arising on nevus spilus. *Int J Dermatol* 2006; 45: 1380–1381.
30. Karam SL, Jackson SM. Malignant melanoma arising within nevus spilus. *Skinmed* 2012; 10: 100–102.
31. Lee SB, Hummel HM, Krengel S, Enk A, Haenssle HA. Mosaic RASopathies: phenotypic and genotypic differentiation of naevus spilus-type congenital melanocytic naevus and segmental naevus spilus. *J Eur Acad Dermatol Venereol* 2016; 30: 1244–1246.
32. Kinsler VA, Krengel S, Riviere JB, Waelchli R, Chapusot C, Al-Olabi L, et al. Next-generation sequencing of nevus spilus-type congenital melanocytic nevus: exquisite genotype-phenotype correlation in mosaic RASopathies. *J Invest Dermatol* 2014; 134: 2658–2660.
33. Haenssle HA, Kaune KM, Buhl T, Thoms KM, Padeken M, Emmerl S, et al. Melanoma arising in segmental nevus spilus: detection by sequential digital dermatoscopy. *J Am Acad Dermatol* 2009; 61: 337–341.
34. Cecchi R, Fancelli L, Troiano M. Melanoma arising in giant zosteriform nevus spilus. *Indian J Dermatol Venereol Leprol* 2012; 78: 643–645.
35. Stella A, Ponholzer K, Weingast J, Binder M. Melanoma arising in a giant nevus spilus maculosus. *Dermatol Pract Concept* 2011; 1: 21–24.
36. Gescheidt-Shoshany H, Weltfriend S, Bergman R. Nodular melanoma arising in a large segmental speckled lentiginous nevus. *Am J Dermatopathol* 2015; 37: 663–664.
37. From L. Congenital nevi – let's be practical. *Pediatr Dermatol* 1992; 9: 345–346.
38. Manganoni AM, Pavoni L, Farisoglio C, Sereni E, Calzavara-Pinton P. Report of 27 cases of naevus spilus in 2134 patients with melanoma: is naevus spilus a risk marker of cutaneous melanoma? *J Eur Acad Dermatol Venereol* 2012; 26: 129–130.
39. Kopf AW, Levine LJ, Rigel DS, Friedman RJ, Levenstein M. Congenital-nevus-like nevi, nevi spili, and cafe-au-lait spots in patients with malignant melanoma. *J Dermatol Surg Oncol* 1985; 11: 275–280.
40. Vidaurri-de la Cruz H, Happel R. Two distinct types of speckled lentiginous nevi characterized by macular versus papular speckles. *Dermatology* 2006; 212: 53–58.
41. Sawicka E, Szczygelski O, Zak K, Peczkowski P, Michalak E, Bekiesinska-Figatowska M. Giant congenital melanocytic nevi: selected aspects of diagnostics and treatment. *Med Sci Monit* 2015; 21: 123–132.