

Table SII. Full coefficients of adjusted hazard ratios (HR) for risk of myocardial infarction in patients with psoriasis

	Any myocardial infarction		First-time myocardial infarction	
	Adjusted HR (95% CI)	p-value	Adjusted HR (95% CI)	p-value
Any psoriasis				
Mild psoriasis	1.02 (0.96–1.09)	0.5209	1.03 (0.96–1.11)	0.3646
Severe psoriasis	1.21 (1.07–1.37)	0.0023	1.21 (1.05–1.39)	0.0065
Sex	2.01 (1.97–1.37)	<0.0001	2.06 (2.02–2.10)	<0.0001
Age	1.06 (1.06–1.06)	<0.0001	1.07 (1.06–1.07)	<0.0001
Socioeconomic status				
Lowest group	1.04 (1.01–1.07)	0.0029	1.03 (1.00–1.06)	0.0814
Below average	1.02 (1.00–1.05)	0.0891	1.02 (0.99–1.05)	0.1778
Average	(reference)		(reference)	
Above average	0.91 (0.88–0.94)	<0.0001	0.91 (0.88–0.95)	<0.0001
Highest group	0.82 (0.80–0.85)	<0.0001	0.82 (0.79–0.84)	<0.0001
Healthcare consumption	1.01 (1.00–1.02)	0.0314	1.01 (1.00–1.02)	0.0250
Previous CVD	2.18 (2.13–2.22)	<0.0001	1.70 (1.66–1.74)	<0.0001
Smoking	2.07 (2.03–2.11)	<0.0001	2.20 (2.15–2.25)	<0.0001
Alcohol abuse	1.25 (1.21–1.30)	<0.0001	1.27 (1.22–1.32)	<0.0001
Diabetes	1.56 (1.52–1.60)	<0.0001	1.63 (1.60–1.68)	<0.0001
Hypertension	1.32 (1.30–1.35)	<0.0001	1.29 (1.27–1.32)	<0.0001
Statin use	0.99 (0.97–1.02)	0.5331	0.90 (0.88–0.92)	<0.0001

CI: confidence interval; CVD: cardiovascular disease.