

**Table SIV. Full coefficients of adjusted hazard ratios (HR) for risk of myocardial infarction in patients with psoriatic arthritis**

	Any myocardial infarction		First-time myocardial infarction	
	Adjusted HR (95% CI)	p-value	Adjusted HR (95% CI)	p-value
<b>Joint disease</b>				
Psoriatic arthritis	1.22 (1.05–1.43)	0.0116	1.23 (1.04–1.47)	0.0177
Sex	2.01 (1.97–2.05)	<0.0001	2.06 (2.02–2.10)	<0.0001
Age	1.06 (1.06–1.06)	<0.0001	1.07 (1.06–1.07)	<0.0001
<b>Socioeconomic status</b>				
Lowest group	1.04 (1.01–1.07)	0.0049	1.03 (1.00–1.06)	0.0904
Below average	1.02 (1.00–1.05)	0.1126	1.02 (0.99–1.05)	0.2138
Average (reference)			(reference)	
Above average	0.91 (0.88–0.94)	<0.0001	0.91 (0.88–0.95)	<0.0001
Highest group	0.82 (0.79–0.85)	<0.0001	0.81 (0.79–0.84)	<0.0001
Healthcare consumption	1.01 (1.00–1.02)	0.0087	1.01 (1.00–1.02)	0.0113
Previous CVD	2.17 (2.13–2.22)	<0.0001	1.70 (1.66–1.74)	<0.0001
Smoking	2.08 (2.04–2.12)	<0.0001	2.21 (2.16–2.26)	<0.0001
Alcohol abuse	1.26 (1.21–1.31)	<0.0001	1.28 (1.23–1.33)	<0.0001
Diabetes	1.55 (1.51–1.60)	<0.0001	1.63 (1.58–1.68)	<0.0001
Hypertension	1.33 (1.30–1.35)	<0.0001	1.30 (1.27–1.33)	<0.0001
Statin use	0.99 (0.97–1.02)	0.5318	0.90 (0.87–0.92)	<0.0001

CI: confidence interval; CVD: cardiovascular disease.