

Table SI. Clinicopathological features of patients with locally aggressive trichoblastic tumours (n = 36)

Clinicopathological features	
Sex, n (%)	
Male	23 (63.9)
Female	13 (36.1)
Median age at diagnosis, years, median (range)	64 (39–98)
Median diameter of the tumour, mm, median (range)	23 (15–50)
Location of the lesions, n (%); MD=1	
Face	21 (60)
Trunk	7 (20)
Limbs	6 (17.1)
Other	1 (2.9)
Median number of excision, n (range)	2 (1–5)
Follow-up, months; MD=16, median (range)	64.2 (10–120)
Invasion depth, n (%)	
Subcutaneous invasion	36 (100)
Muscle invasion	18 (50)
Bland round to ovoid finely nucleolated nucleus, n (%)	36 (100)
Presence of mitosis, n (%)	31 (86.1)
Apoptotic bodies, n (%)	31 (86.1)
Follicular stroma, n (%)	36 (100)
Focal connection with epidermis, n (%)	12 (33.3)
Focal microscopic ulceration, n (%)	6 (16.7)
Poor circumscription with infiltrative borders, n (%)	36 (100)
Retraction cleft between stroma and adjacent connective tissue, n (%)	5 (13.9)
Calcification, n (%)	24 (66.6)
Perineural infiltration, n (%)	3 (8.3)
Vascular invasion, n (%)	–

MD: missing data.