Appendix S1.

For the second part of the HS questionnaire, we included 3 items with photographs (items 5, 6, 7) followed by 3 items in accordance with the definition of HS by the HS Foundation (12). The photographs reflected the body area involved in HS: armpit (Q. 5), inguinal creases (Q. 6), bottom (Q. 7). There were 2–4 photographs per item, including different elementary lesions such as abscesses or scars observed in Hurley 1 and 2 patients. The Hurley score is a severity score ranging from 1 to 3 (13). We did not include Hurley 3 score photographs, as it is less common than the other scores (14).

For the second part of the psoriasis questionnaire, we included 3 items with photographs (items 5, 6, 7) followed by 2 items regarding the presence of inverse psoriasis or a possible differential diagnosis (hives or eczema). These items were derived from expert opinion and the National Psoriasis Foundation's survey on psoriasis (8). The pictures reflected the body area involved in psoriasis: elbows, lumbosacral (Q. 5), nails (Q. 6), scalp (Q. 7). There were 1–2 photographs per item, including typical lesions, such as erythemato-squamous patches and punctuate depressions in the nail plate.

For the second part of the vitiligo questionnaire, we included one item with a panel of photographs (Q. 8) and 3 items which focused on typical areas involved in vitiligo and common presentations. The photographs reflected typical symmetrical macules of depigmented skin.