

Table SII. Anaesthesia methods and botulinum toxin (BTX) treatment

| | Palms | Axillae | Feet | Face/head | Back | Groin | Multifocal |
|-----------------------------------------------|-------|---------|-------|-----------|-------|-------|------------|
| Anaesthesia (number of patients, <i>n</i>) | | | | | | | |
| Topical | | 25 | | | | 1 | 1 |
| IVRA | 30 | | 2 | | | | 8 |
| Regional | 31 | | | | | | 12 |
| Alfentanil | 3 | 1 | | 6 | 1 | 1 | 15 |
| General anaesthesia | 5 | | 2 | | | | 5 |
| Ice | | 2 | | | | | |
| Combination of anaesthesia | 34 | 1 | 4 | | | | 63 |
| No anaesthesia | | 61 | | 1 | | | |
| BTX treatment (number of patients, <i>n</i>) | | | | | | | |
| Dysport | | 80 | | | | | |
| Xeomin | | 3 | 1 | | | 1 | |
| NeuroBloc | 2 | 3 | | 7 | 1 | | 5 |
| Dysport + NeuroBloc | 93 | 4 | 7 | | | 1 | 91 |
| Xeomin + NeuroBloc | 8 | | | | | | 8 |
| BTX dose (median units) | | | | | | | |
| Dysport | | 300 | | | | | |
| Xeomin | | 100 | 310 | | | 180 | |
| NeuroBloc | 2,563 | 350 | | 1,625 | 3,000 | | 3,625 |
| Dysport (D) + NeuroBloc (N) | | | | | | | |
| | D 500 | D 200 | D 600 | | | D 350 | D 600 |
| | N 250 | N 500 | N 875 | | | N 875 | N 750 |
| Xeomin (X) + NeuroBloc (N) | | | | | | | |
| | X 210 | | | | | | X 278 |
| | N 250 | | | | | | N 250 |

IVRA: intravenous regional anaesthesia; BTX: botulinum toxin.