

Table SI. Baseline characteristics

Characteristics	Total	SE	PDT	5-FU	Other*	p-value
<i>Patients characteristics</i>						
Sex, n (%)	n = 608	n = 241	n = 296	n = 46	n = 25	
Female	347 (57.1)	122 (50.6)	190 (64.2)	23 (50)	12 (48)	0.007**
Male	261 (42.9)	119 (49.4)	106 (35.8)	23 (50)	13 (52)	
Age at diagnosis, years, mean (SD)	73 (10.8)	75 (9.9)	72 (11.0)	70 (12.1)	73 (13.0)	0.010***
<i>Immunosuppressive medication, n (%)</i>						
No	535 (88.0)	210 (87.1)	262 (88.5)	41 (89.1)	22 (88)	0.95**
Yes	72 (11.8)	30 (12.4)	34 (11.5)	5 (10.9)	3 (12)	
Unknown	1 (0.2)	1 (0.4)	0	0	0	
<i>Tumour characteristics</i>						
Tumour location, n (%)	n = 841	n = 288	n = 450	n = 72	n = 31	
Head neck region	298 (35.4)	102 (35.4)	157 (34.9)	21 (29.2)	18 (58.1)	<0.001**
Ear	32 (3.8)	13 (4.5)	8 (1.8)	6 (8.3)	5 (16.1)	
Upper extremities	164 (19.5)	70 (24.3)	72 (16.0)	18 (25.0)	4 (12.9)	
Lower extremities	205 (24.4)	53 (18.4)	138 (30.7)	14 (19.4)	0	
Trunk	142 (16.9)	50 (17.4)	75 (16.7)	13 (18.1)	4 (12.9)	
<i>Tumour diameter</i>						
≤ 10 mm	497 (59.1)	181 (62.8)	254 (56.4)	41 (56.9)	21 (67.7)	0.51**
11–≤ 30 mm	240 (28.5)	75 (26.0)	138 (30.7)	22 (30.6)	5 (16.1)	
> 30 mm	10 (1.2)	3 (1.0)	6 (1.3)	1 (1.4)	0	
Unknown	94 (11.2)	29 (10.1)	52 (11.6)	8 (11.1)	5 (16.1)	

*Cryotherapy, imiquimod or curettage. **Group differences were tested with χ^2 test. ***Group differences were tested with the one-way analysis of variance (ANOVA).

SE: surgical excision; PDT: photodynamic therapy; 5-FU: 5-fluorouracil; 95% CI: 95% confidence interval.