

Table SI. Characteristics of patients with cutaneous chronic graft versus host disease (GVHD) (*n* = 20)

No.	Sex/age at transplantation, years	Donor and diagnosis graft source	Type of cutaneous CGVHD	Extent of cutaneous CGVHD	Cutaneous features of lichenoid to sclerotic rash, months	Time interval	Additional dermatological findings	Therapy	Cutaneous response/ Systemic therapy
1	M/5	OP	MUD, BM,	<i>De novo</i>	Localized to photo-exposed areas	Lichenoid	–	Photo-distribution of rash	Topical
2	F/7	AML	MUD, BM	<i>De novo</i>	Localized	Lichenoid	–	Topical	Yes
3	F/4	ALCL	MSD, BM	Progressive	Widespread	Lichenoid and then sclerotic	Keratoderma	Topical, UVA1	Yes
4	M/10	CML	MUD, BM	Progressive	Widespread	Lichenoid and then sclerotic	Keratoderma, scarring alopecia	Topical, NB UVB, UVA1, ECP, systemic	No/prednisone, tacrolimus
5 ^a	F/15	AML	MSD, BM	<i>De novo</i>	Localized to the scalp	Lichenoid	–	Topical, systemic	Death
6	M/0	HES	MSD, BM	<i>De novo</i>	Localized to the scalp	Lichenoid	–	Topical	Yes
7	F/7	HLH	MUD, BM	Progressive	Widespread	Lichenoid and then sclerotic	Scarring alopecia	Topical, NB UVB, UVA1, ECP, systemic	No/prednisone, IV lidocaine, MTX
8	M/14	ALL	MUD, BM	Progressive	Widespread	Lichenoid and then sclerotic	Keratoderma	Topical, ECP, systemic	Yes/prednisone, MMF, tacrolimus
9	M/15	ALL	MSD, BM	<i>De novo</i>	Localized to the face and upper trunk	Atopic dermatitis-like	–	Topical	Yes
10	M/3	AML	MRD, BM	<i>De novo</i>	Widespread	Lichenoid and then sclerotic	Keratoderma, scarring alopecia	Topical, NB UVB, systemic	No/prednisone
11 ^a	M/8	NB	MUD, PBSC	Progressive	Widespread	Lichenoid	–	Systemic	Death
12	M/7	ALL	MUD, PBSC	Progressive	Localized to the face and upper trunk	Lichenoid	–	Topical, systemic	Rejection of graft
13	M/19	AML	MSD, PBSC	<i>De novo</i>	Widespread	Lichenoid and then sclerotic, erosions	–	Topical, ECP, systemic	b
14	M/10	ALL	MSD, PBSC	Progressive	Widespread	Lichenoid and then sclerotic, erosions	Scarring alopecia	Topical, ECP, systemic	No/prednisone, cyclosporine, MMF, imatinib, MSC
15	M/18	ALL	MSD, PBSC	<i>De novo</i>	N/A	Lichenoid	–	ECP, systemic	b
16 ^a	M/11	ALL	MSD, PBSC	<i>De novo</i>	Widespread	Lichenoid	–	Topical, ECP, systemic	Death
17	M/18	ALL	MSD, PBSC	<i>De novo</i>	Localized to the face and upper chest	Lichenoid	–	Photo-distribution of rash	b
18	F/5	ALL	MSD, PBSC	<i>De novo</i>	Widespread	Lichenoid	–	Topical, systemic	LFU
19	F/6	ALL	MUD, PBSC	Progressive	Localized to the limbs	Lichenoid	–	Topical, ECP, systemic	Yes
20	F/8	ALL	MSD, PBSC	Progressive	Localized to the limbs, trunk and scalp	Ichthyosiform	–	Topical, ECP, systemic	Yes

^aThese patients died. ^bIn these patients systemic therapies were required for involvement of other organs, such as gastrointestinal tract, liver and lungs, and their cutaneous disease was mild or less clinically relevant: ALCL: anaplastic large cell lymphoma; ALL: acute lymphoblastic leukaemia; AML: acute myeloid leukaemia; BM: bone marrow; cGVHD: chronic graft versus host disease; CMV: chronic myeloid leukaemia; ECP: extracorporeal photopheresis; HES: hypereosinophilic syndrome; HLH: haemophagocytic lymphohistiocytosis; IV: intravenous; LFU: loss of follow-up; MRD: HLA-matched non-sibling relative donor; MMF: mycophenolic acid; MSC: mesenchymal stem cells; MSD: HLA-matched sibling donor; MTX: methotrexate; MUD: HLA-matched unrelated donor; NB: non-applicable; N/A: non-applicable; NB: neuroblastoma; NB UVB: narrow-band ultraviolet B; OP: osteopetrosis; PBSC: peripheral blood stem cell; UVA1: ultraviolet A1.