

Table SI. Clinical variables evaluated

Demographic data	Age (years), sex, profession
Predominant type of urticaria	Acute urticaria, chronic urticaria, chronic inducible urticaria
Presence of angioedema	
Concomitant CIndU	Demographic urticaria, delay pressure urticaria, cholinergic urticaria, cold contact urticaria, heat contact urticaria, solar urticaria, vibratory urticaria, AQuagenic urticaria, contact urticaria, exercise-induced anaphylaxis/urticaria
Disease history	Total disease duration (months), number of episodes during the patient's life, mean time until the first visit (months), duration of current episode (months)
Disease activity	Urticaria Activity Score, Urticaria Activity Score 7, Urticaria Activity Score 3 weeks
Personal history and comorbidities	Drug and alcohol intake, atopy, urticaria, thyroid diseases, rheumatological and autoimmune disorders, infections, digestive system disorders, psychiatric diseases, diabetes mellitus/endocrine disorders, hypertension and other renal disorders, cardiovascular diseases, gynaecological diseases, haematological diseases, genetic disorders, other dermatological diseases, benign and malignant neoplasms
Triggering/exacerbating factors	Infections, stress, insect bite, drugs prescribed drugs, food, systemic disease

CIndU: concomitant inducible urticaria.