

Table SII. Laboratory parameters and other diagnostic tests

Laboratory workup (units) [normal values range]	Other tests
Cell blood count	ASST
Kidney function test	APST
Urea (mg/dl) [10–50] Creatinine (mg/dl) [0.5–1.2] Glomerular filtration rate (ml/min/1.73 m ²) [>60]	
Liver function test	Basophil Activation Test
Bilirubin (mg/dl) [0.2–1.2] Liver enzymes (ALT (UI/l 37C) [7–31], AST (UI/l 37C) [10–32] Gamma glutamyl transpeptidase (GGT) (UI/l 37C) [8–61] Phosphatase alkaline (UI/l 37C) [35–104]	
Lipid profile	Skin biopsy
Cholesterol (mg/dl) [120–200]	
Acute-phase reactants	Parasites in faeces
Erythrocyte sedimentation rate (mm/h) [0–12] C-reactive protein (mg/dl) [0–0.8] Ferritin (ng/ml) [13–150] Rheumatoid factor (U/ml) [0.01–20]	
Thyroid function	Helicobacter pylori breath test
Thyroid hormone test (triiodothyronine (pg/ml) [2.0–4.4] Thyroxin (ng/dl) [0.72–1.7] Thyroid-stimulating hormone (mcU/ml) [0.35–5] Anti-thyroid peroxidase antibodies (UI/ml) [0.1–35] Anti-thyroglobulin antibodies (UI/ml) [0.1–40]	
Other laboratory parameters	Patch test
IgE (kU/l) [0–100] D-dimer (ng/ml) [0–150]	
Immunologic profile	Specific IgE
Antinuclear antibodies Anti-DNA ds (UI/ml) [0.1–300] Complement system (C4 (mg/dl) [10–40], CH50 (U/ml))	

ASST: Autologous Serum Skin Test; APST: Autologous Plasma Skin Test.